

Policy Number	SUR703.032
Policy Effective Date	05/15/2026

Allogeneic Hematopoietic Cell Transplantation for Myelodysplastic Syndromes and Myeloproliferative Neoplasms

Table of Contents
Coverage
Policy Guidelines
Description
Rationale
Coding
References
Policy History

Related Policies (if applicable)
None

Disclaimer

Carefully check state regulations and/or the member contract.

Each benefit plan, summary plan description or contract defines which services are covered, which services are excluded, and which services are subject to dollar caps or other limitations, conditions or exclusions. Members and their providers have the responsibility for consulting the member's benefit plan, summary plan description or contract to determine if there are any exclusions or other benefit limitations applicable to this service or supply. **If there is a discrepancy between a Medical Policy and a member's benefit plan, summary plan description or contract, the benefit plan, summary plan description or contract will govern.**

Coverage

Allogeneic hematopoietic cell transplantation **may be considered medically necessary** as a treatment of:

- Myelodysplastic syndromes (see Policy Guidelines section); or
- Myeloproliferative neoplasms (see Policy Guidelines section).

Allogeneic hematopoietic cell transplantation for myelodysplastic syndromes and myeloproliferative neoplasms that do not meet the criteria in the Policy Guidelines section **is considered experimental, investigational and/or unproven.**

Policy Guidelines

Myeloid Neoplasms

Myeloid neoplasms are categorized according to criteria developed by the World Health Organization (WHO). Neoplasms are risk-stratified using the International Prognostic Scoring System (IPSS).

2022 WHO Classification Scheme for Myeloid Neoplasm and Histiocytic/Dendritic Neoplasms

Clonal hematopoiesis (CH)

- CH of indeterminate potential (CHIP)
- Clonal cytopenia of undetermined significance (CCUS)

Myeloproliferative neoplasms (MPN)

- Chronic myeloid leukemia (CML), BCR-ABL1⁺
- Chronic neutrophilic leukemia (CNL)
- Polycythemia vera
- Primary myelofibrosis (PMF)
- Essential thrombocythemia
- Chronic eosinophilic leukemia
- MPN, not otherwise specified
- Juvenile myelomonocytic leukemia

Mastocytosis

- Cutaneous mastocytosis
- Systemic mastocytosis
- Mast cell sarcoma

Childhood myelodysplastic syndromes (MDS)

- Childhood MDS with low blasts
 - Hypocellular
 - Not otherwise specified
- Childhood MDS with increased blasts

Myeloid/lymphoid neoplasms with eosinophilia and tyrosine kinase gene fusions (MLN-TK)

Myelodysplastic/myeloproliferative neoplasms (MDS/MPN)

- Chronic myelomonocytic leukemia (CMML)
- MDS/MPN with neutrophilia
- MDS/MPN with *SF3B1* mutation and thrombocytosis
- MDS/MPN, not otherwise specified

Myelodysplastic neoplasms (MDS)

- MDS with defining genetic abnormalities
 - MDS with low blasts and isolated 5q deletion (MDS-5q)
 - MDS with low blasts and *SF3B1* mutation (MDS-*SF3B1*), or MDS with low blasts and ring sideroblasts
 - MDS with biallelic *TP53* inactivation (MDS-bi*TP53*)

- MDS, morphologically defined
 - MDS with low blasts (MDS-LB)
 - MDS, hypoplastic (MDS-h)
 - MDs with increased blasts (MDS-IB)
 - MDS-IB1
 - MDS-IB2
 - MDS with fibrosis (MDS-f)

Acute myeloid leukemia (AML)

- AML with defining genetic abnormalities
- AML, defined by differentiation

Secondary myeloid neoplasms

- Myeloid neoplasms post cytotoxic therapy
- Myeloid neoplasms associated with germline predisposition

Dendritic cell and histiocytic neoplasms

- Plasmacytoid dendritic cell neoplasms
- Langerhans cell and other dendritic cell neoplasms
- Histiocytic neoplasms

Acute leukemias of ambiguous lineage (ALAL)

- ALAL with defining genetic abnormalities
- ALAL, immunophenotypically defined

Genetic tumor syndromes with predisposition to myeloid neoplasia

Risk Stratification of Myelodysplastic Syndromes

Risk stratification for MDS is performed using the IPSS (Table PG1). This system was developed after pooling data from 7 studies that used independent, risk-based prognostic factors. The prognostic model and the scoring system were based on blast count, degree of cytopenia, and blast percentage. Risk scores were weighted relative to their statistical power. This system is widely used to group individuals into either low-risk or high-risk groups (Table PG2). The low-risk group includes low-risk and intermediate-1 IPSS groups; treatment goals in low-risk MDS individuals are to improve quality of life and achieve transfusion independence. In the high-risk group, which includes intermediate-2 and high-risk IPSS groups, treatment goals are slowing disease progression to AML and improving survival. IPSS is usually calculated on diagnosis. The role of lactate dehydrogenase, marrow fibrosis, and β_2 -microglobulin also should be considered after establishing IPSS. If elevated, the prognostic category worsens by 1 category change.

Table PG1. International Prognostic Scoring System: Myelodysplastic Syndrome Prognostic Variables

Variable	0	0.5	1.0	1.5	2.0
----------	---	-----	-----	-----	-----

Marrow blasts, %	<5	5 to 10	NA	11 to 20	21 to 30
Karyotype	Good	Intermediate	Poor	NA	NA
Cytopenias	0/1	2/3	NA	NA	NA

NA: not applicable.

Table PG2. International Prognostic Scoring System: Myelodysplastic Syndromes Clinical Outcomes

Risk Group	Total Score	Median Survival, years	Time for 25% of patients to Progress to AML
Low	0	5.7	9.4 years
Intermediate-1	0.5 to 1.0	3.5	3.3 years
Intermediate-2	1.5 to 2.0	1.2	1.12 years
High	≥2.5	0.4	0.2 years

AML: acute myelocytic leukemia.

An updated 5-category IPSS has been proposed for prognosis in individuals with primary MDS or secondary AML to account for chromosomal abnormalities frequently seen in MDS (see Schanz et al. 2012). This system stratifies patients into 5 categories: very poor, poor, intermediate, good, and very good. There has also been an investigation into using the 5-category IPSS to better characterize risk in MDS.

Given the long natural history of MDS, allogeneic hematopoietic cell transplantation (allo-HCT) is typically considered in individuals with increasing numbers of blasts, signaling a possible transformation to AML. Subtypes falling into this category include refractory anemia with excess blasts, refractory anemia with excess blasts in transformation, or CMML.

Individuals with refractory anemia with or without ringed sideroblasts may be considered candidates for allo-HCT when chromosomal abnormalities are present, or when the disorder is associated with the development of significant cytopenias (e.g., neutrophils <500/mm³, platelets <20,000/mm³).

Individuals with myeloproliferative neoplasms may be considered candidates for allo-HCT when there is a progression to myelofibrosis or toward acute leukemia. In addition, allo-HCT may be considered in individuals with essential thrombocythemia with an associated thrombotic or hemorrhagic disorder. Use of allo-HCT should be based on the following criteria: cytopenias, transfusion dependence, increasing blast percentage over 5%, and age.

Some individuals for whom a conventional myeloablative allo-HCT could be curative may be candidates for reduced-intensity conditioning (RIC) allo-HCT. These include individuals whose age (typically >60 years) or comorbidities (e.g., liver or kidney dysfunction, generalized debilitation, prior intensive chemotherapy, low Karnofsky Performance Status) preclude the use of a standard myeloablative conditioning (MAC) regimen. The ideal allogeneic donors

are human leukocyte antigen (HLA)-identical siblings, matched at the HLA-A, -B, and -DR loci (6/6). Related donors mismatched at 1 locus are also considered suitable donors. A matched, unrelated donor identified through the National Marrow Donor Registry is typically the next option considered. Recently, there has been interest in haploidentical donors, typically a parent or a child of the individual, who usually share only 3 of the 6 major histocompatibility antigens. Most individuals will have such a donor; however, the risk of graft-versus-host disease (GVHD) and overall morbidity of the procedure may be severe, and experience with these donors is not as GVHD extensive as that with matched donors.

Evidence and clinical guidelines suggest RIC allo-HCT may be considered as a risk-adapted strategy for high-risk individuals of MAC-intolerance as follows:

MDS

- Older age
- IPSS intermediate-2 or high risk
- Multiple comorbidities (e.g., hematopoietic cell transplantation-comorbidity index [HCT-CI] score higher than 2)
- Red blood cell transfusion dependence
- Neutropenia
- Thrombocytopenia
- High-risk cytogenetics
- Increasing blast percentage

Myeloproliferative neoplasm

- Cytopenias
- Transfusion dependence
- Increasing blast percentage over 5%
- Age 60 to 65 years

Description

Myelodysplastic syndromes and myeloproliferative neoplasms refer to a heterogeneous group of clonal hematopoietic disorders with the potential to transform into acute myelocytic leukemia. Allogeneic hematopoietic cell transplantation (allo-HCT) has been proposed as a curative treatment option for patients with these disorders.

Myelodysplastic Syndromes

Myelodysplastic syndrome (MDS) can occur as a primary (idiopathic) disease or can be secondary to cytotoxic therapy, ionizing radiation, or other environmental insults. Chromosomal abnormalities are seen in 40% to 60% of patients, frequently involving deletions of chromosome 5 or 7 or an extra chromosome as in trisomy 8. Most MDS diagnoses occur in individuals older than age 55 to 60 years, with an age-adjusted incidence of 62% among individuals older than age 70 years. Patients succumb either to disease progression to acute

myeloid leukemia (AML) or to complications of pancytopenias. Patients with higher blast counts or complex cytogenetic abnormalities have a greater likelihood of progressing to AML than do other patients.

Myelodysplastic Syndrome Classification and Prognosis

The French-American-British system was previously used to classify MDS into 5 subtypes: 1) refractory anemia; 2) refractory anemia with ringed sideroblasts; 3) refractory anemia with excess blasts; 4) refractory anemia with excess blasts in transformation; and 5) chronic myelomonocytic leukemia. The French-American-British system was supplanted by that of the World Health Organization (WHO), which differentiates between MDS defined by genetic abnormalities or by morphologic features (in the form of dysplastic cell lineages), and reduces the threshold maximum blast percentage for the diagnosis of MDS from 30% to 20%. (1)

The most commonly used prognostic scoring system for MDS is the International Prognostic Scoring System (IPSS), which groups patients into 1 of 4 prognostic categories based on the number of cytopenias, cytogenetic profile, and the percentage of blasts in the bone marrow. This system underweights the clinical importance of severe, life-threatening neutropenia and thrombocytopenia in therapeutic decisions and does not account for the rate of change in critical parameters (e.g., peripheral blood counts, blast percentage). However, the IPSS has been useful in a comparative analysis of clinical trial results, and its utility confirmed at many institutions. An updated 5-category IPSS has been proposed for prognosis in patients with primary MDS or secondary AML to account for chromosomal abnormalities frequently seen in MDS. (2) This system stratifies patients into 5 categories: very poor, poor, intermediate, good, and very good. There has been an investigation into using the 5-category IPSS to better characterize risk in MDS. A second prognostic scoring system incorporates the WHO subgroup classification that accounts for blast percentage, cytogenetics, and severity of cytopenias as assessed by transfusion requirements. The WHO classification-based Prognostic Scoring System uses a 6-category system, which allows more precise prognostication of overall survival (OS) duration, as well as risk for progression to AML.

Myelodysplastic Syndrome Treatment

Treatment of nonprogressing MDS has previously involved best supportive care, including red blood cell and platelet transfusions and antibiotics. Active therapy was given only when MDS progressed to AML or resembled AML with severe cytopenias. An array of therapies are now available to treat MDS, including hematopoietic growth factors (e.g., erythropoietin, darbepoetin, granulocyte colony-stimulating factor), transcriptional-modifying therapy (e.g., U.S. Food and Drug Administration [FDA] approved hypomethylating agents, nonapproved histone deacetylase inhibitors), immunomodulators (e.g., lenalidomide, thalidomide, antithymocyte globulin, cyclosporine A), low-dose chemotherapy (e.g., cytarabine), and allo-HCT. Given the spectrum of treatments available, the goal of therapy must be decided upfront whether it is to improve anemia, thrombocytopenia, or neutropenia, to eliminate the need for red blood cell transfusion, to achieve complete remission, or to cure the disease.

Allo-HCT is the only approach with curative potential, but its use is governed by patient age, performance status, medical comorbidities, the patient's preference, risk category, and severity of MDS at presentation. Allo-HCT is discussed in more detail in a subsequent section.

Chronic Myeloproliferative Neoplasms

Chronic myeloproliferative neoplasms are clonal bone marrow stem cell disorders; as a group, approximately 8,400 myeloproliferative neoplasms are diagnosed annually in the United States. Like MDS, myeloproliferative neoplasms primarily occur in older individuals, with approximately 67% reported in patients aged 60 years and older.

Myeloproliferative neoplasms are characterized by the slow but progressive expansion of a clone of cells with the potential evolution into a blast crisis similar to AML. Myeloproliferative neoplasms share a common stem cell-derived clonal heritage, with phenotypic diversity attributed to abnormal variations in signal transduction as the result of a spectrum of variants that affects protein tyrosine kinases or related molecules. The unifying characteristic common to all myeloproliferative neoplasms is effective clonal myeloproliferation resulting in peripheral granulocytosis, thrombocytosis, or erythrocytosis that is devoid of dyserythropoiesis, granulocytic dysplasia, or monocytosis.

Myeloproliferative Neoplasm Classification

Myeloproliferative neoplasms are a subdivision of myeloid neoplasms that includes 4 classic disorders: chronic myeloid leukemia, polycythemia vera, essential thrombocytopenia, and primary myelofibrosis. The WHO classification also includes chronic neutrophilic leukemia, chronic eosinophilic leukemia not otherwise specified, and myeloproliferative neoplasm unclassifiable. In the 2016 classification, mastocytosis is no longer considered a subgroup of the myeloproliferative neoplasms due to its unique clinical and pathologic features.

Myeloproliferative Neoplasm Treatment

In indolent, nonprogressing cases, therapeutic approaches are based on relief of symptoms. Supportive therapy may include prevention of thromboembolic events. Hydroxyurea may be used in cases of high-risk essential thrombocytosis and polycythemia vera, and intermediate- and high-risk primary myelofibrosis.

The FDA (2011) approved the orally administered selective Janus kinase 1 and 2 inhibitor ruxolitinib for the treatment of intermediate- or high-risk myelofibrosis. Ruxolitinib has been associated with improved OS, spleen size, and symptoms of myelofibrosis compared with placebo. (3) The Randomized Study of Ruxolitinib Tablets Compared to Best Available Therapy in Subjects With Primary Myelofibrosis, Post-Polycythemia Vera-Myelofibrosis or Post-Essential Thrombocythemia Myelofibrosis (COMFORT-II trial [2013]) compared ruxolitinib with best available therapy in patients who had intermediate- and high-risk myelofibrosis, and demonstrated improvements in spleen volume and OS. (4) In a randomized trial comparing ruxolitinib with best available therapy (including antineoplastic agents, most commonly hydroxyurea, glucocorticoids) with no therapy for treatment of myelofibrosis, Harrison et al. (2012) reported improvements in spleen size and quality of life, but not OS. (5) In 2019, the

FDA also approved fedratinib (Inrebic®) for adults with intermediate-2 or high-risk primary or secondary myelofibrosis based on results from a double-blind, randomized, placebo-controlled trial that found improvement in spleen volume and myelofibrosis-related symptoms. (6)

Myeloablative allo-HCT has been considered the only potentially curative therapy, but because most patients are of advanced age with attendant comorbidities, its use is limited to those who can tolerate the often-severe treatment-related adverse events of this procedure. However, the use of reduced-intensity conditioning (RIC) for allo-HCT has extended the potential benefits of this procedure to selected individuals with these disorders. Allo-HCT is discussed in more detail in the next section.

Hematopoietic Cell Transplantation

Hematopoietic cell transplantation (HCT) is a procedure in which hematopoietic stem cells are intravenously infused to restore bone marrow and immune function in cancer patients who receive bone marrow-toxic doses of cytotoxic drugs with or without whole-body radiotherapy. Hematopoietic stem cells may be obtained from the transplant recipient (autologous HCT) or a donor (allo-HCT). They can be harvested from bone marrow, peripheral blood, or umbilical cord blood shortly after delivery of neonates.

Immunologic compatibility between infused hematopoietic stem cells and the recipient is not an issue in autologous HCT. In allogeneic stem cell transplantation, immunologic compatibility between donor and patient is a critical factor for achieving a successful outcome. Compatibility is established by typing of human leukocyte antigens (HLA) using cellular, serologic, or molecular techniques. Human leukocyte antigen refers to the gene complex expressed at the HLA-A, -B, and -DR (antigen-D related) loci on each arm of chromosome 6. An acceptable donor will match the patient at all or most of the HLA loci.

Conditioning for Hematopoietic Cell Transplantation

Conventional Conditioning

The conventional (“classical”) practice of allo-HCT involves administration of cytotoxic agents (e.g., cyclophosphamide, busulfan) with or without total body irradiation at doses sufficient to cause bone marrow ablation in the recipient. The beneficial treatment effect of this procedure is due to a combination of the initial eradication of malignant cells and subsequent graft-versus-malignancy effect mediated by non-self-immunologic effector cells. While the slower graft-versus-malignancy effect is considered the potentially curative component, it may be overwhelmed by existing disease in the absence of pretransplant conditioning. Intense conditioning regimens are limited to patients who are sufficiently medically fit to tolerate substantial adverse effects. These include opportunistic infections secondary to loss of endogenous bone marrow function and organ damage or failure caused by cytotoxic drugs. Subsequent to graft infusion in allo-HCT, immunosuppressant drugs are required to minimize graft rejection and graft-versus-host disease (GVHD), which increases susceptibility to opportunistic infections.

The success of autologous HCT is predicated on the potential of cytotoxic chemotherapy, with or without radiotherapy, to eradicate cancerous cells from the blood and bone marrow. This permits subsequent engraftment and repopulation of the bone marrow with presumably normal hematopoietic stem cells obtained from the patient before undergoing bone marrow ablation. Therefore, autologous HCT is typically performed as consolidation therapy when the patient's disease is in complete remission. Patients who undergo autologous HCT are also susceptible to chemotherapy-related toxicities and opportunistic infections before engraftment, but not GVHD.

Reduced-Intensity Conditioning Allogeneic Hematopoietic Cell Transplantation

Reduced-intensity conditioning refers to the pretransplant use of lower doses of cytotoxic drugs or less intense regimens of radiotherapy than are used in traditional full-dose myeloablative conditioning (MAC) treatments. Although the definition of RIC is variable, with numerous versions employed, all regimens seek to balance the competing effects of relapse due to residual disease and non-relapse mortality. The goal of RIC is to reduce disease burden and to minimize associated treatment-related morbidity and non-relapse mortality in the period during which the beneficial graft-versus-malignancy effect of allogeneic transplantation develops. RIC regimens range from nearly total myeloablative to minimally myeloablative with lymphoablation, with intensity tailored to specific diseases and patient condition. Patients who undergo RIC with allo-HCT initially demonstrate donor cell engraftment and bone marrow mixed chimerism. Most will subsequently convert to full-donor chimerism. In this policy, the term RIC will refer to all conditioning regimens intended to be nonmyeloablative.

Regulatory Status

The U.S. Food and Drug Administration regulates human cells and tissues intended for implantation, transplantation, or infusion through the Center for Biologics Evaluation and Research, under Code of Federal Regulation (CFR) Title 21, parts 1270 and 1271. Hematopoietic stem cells are included in these regulations.

Rationale

Medical policies assess the clinical evidence to determine whether the use of technology improves the net health outcome. Broadly defined, health outcomes are the length of life, quality of life, and ability to function including benefits and harms. Every clinical condition has specific outcomes that are important to patients and to managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of technology, 2 domains are examined: the relevance, and the quality and credibility. To be relevant, studies must represent 1 or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable

intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. Randomized controlled trials are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

Myelodysplastic Syndromes

Clinical Context and Therapy Purpose

The purpose of myeloablative conditioning (MAC) or reduced-intensity conditioning (RIC) allogeneic hematopoietic cell transplant (allo-HCT) in individuals who have myelodysplastic syndrome (MDS) is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this policy.

Populations

The relevant population of interest is individuals with MDS.

Interventions

The therapies being considered are myeloablative or RIC allo-HCT.

Comparators

The following therapies are currently being used: standard of care.

Outcomes

The general outcomes of interest are mortality and morbidity. Beneficial outcomes are an improvement in overall survival (OS) and disease-specific survival. Harmful outcomes are treatment-related morbidity and mortality. Follow-up over months to years is of interest for relevant outcomes.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs.
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Myeloablative Conditioning Allogeneic Hematopoietic Cell Transplantation

Despite the successes seen with drugs now available to treat MDS (e.g., decitabine, azacitidine, lenalidomide), allo-HCT is the only treatment capable of complete and permanent eradication of the MDS clone. (7)

Systematic Reviews

A 2009 review of HCT for MDS evaluated the evidence for allo-HCT with myeloablative conditioning (MAC) for MDS. (8) Reviewers selected 24 studies (prospective and retrospective) published between 2000 and 2008 that included a total 1378 cases (age range, 32 to 59 years). Most patients (n=885) received matched-related donor allo-HCT, with other donor types including syngeneic, matched, unrelated donor, mismatched unrelated donor, and umbilical cord blood. Most studies included de novo and secondary MDS, chronic myelomonocytic leukemia, myeloproliferative neoplasms, de novo and secondary acute myeloid leukemia (AML), and transformed AML. Peripheral blood and bone marrow stem cell grafts were allowed in most studies. The most commonly used conditioning regimens were busulfan plus cyclophosphamide (CY) and CY plus total body irradiation, with cyclosporine A used for graft-versus-host disease (GVHD) prophylaxis. Length of follow-up ranged from 5 months to approximately 8 years. Acute GVHD (grades II to IV) varied from 18% to 100%. Relapse risk ranged from 24% at 1 year to 36% at 5 years. The OS rates ranged from 25% at 2 years to 52% at 4 years, with nonrelapse mortality ranging from 19% at day 100 to 61% at 5 years.

A 2009 review from the American Society for Blood and Marrow Transplantation evaluated the evidence related to HCT in the therapy of MDS, with associated treatment recommendations. (9) Reviewers concluded that outcomes improved with early HCT for patients with an International Prognostic Scoring System (IPSS) score of intermediate-2 or high-risk at diagnosis who had a suitable donor and met the transplant center's eligibility criteria, and for selected patients with a low or intermediate-1 risk IPSS score at diagnosis who had a poor prognostic feature not included in the IPSS (i.e., older age, refractory cytopenias). Koenecke et al. (2015) evaluated the impact on the revised 5-category IPSS score (IPSS-5) on outcomes after HCT in patients with MDS or secondary AML (evolved from MDS). (10) In a cohort of 903 patients retrospectively identified from the European Society for Blood and Marrow Transplantation database, those with poor and very poor risk had shorter relapse-free survival (RFS) and OS than those with very good, good, or intermediate risk. However, the ways that transplant management strategies should change based on cytogenetic abnormalities are not currently well defined.

Reduced-Intensity Conditioning Allogeneic Hematopoietic Cell Transplantation

Systematic Reviews

Song et al. (2021) evaluated the efficacy of RIC followed by allo-HCT in patients with AML and MDS via a meta-analysis of 6 RCTs (N=1413). (11) The 6 RCTs compared RIC to MAC before first allo-HCT in patients with AML in complete remission or MDS, had a median follow-up of >1 year, and displayed a low risk of bias. The primary endpoint was OS. Results revealed that OS was not significantly different between RIC and MAC (hazard ratio [HR], 0.95; 95% confidence interval [CI], 0.64 to 1.4; p=.80), with combined long-term follow-up data also showing no difference in OS between the 2 conditioning approaches (HR, 0.86; 95% CI, 0.53 to 1.41; p=.56).

The cumulative incidence of relapse was also similar between the groups (HR, 1.18; 95% CI, 0.88 to 1.49; $p=.28$). Nonrelapse mortality was significantly improved with RIC as compared to total body irradiation/busulfan-based MAC (HR, 0.53; 95% CI, 0.36 to 0.8; $p=.002$); however, treosulfan-based MAC significantly reduced nonrelapse mortality as compared to RIC (HR, 1.67; 95% CI, 1.02 to 2.72; $p=.04$). RIC was associated with a trend of increasing graft failure ($p=.06$); however, graft failure in both arms was rare. The median duration of follow-up among the studies ranged from 12 to 119 months. The authors concluded that RIC is recommended as an adequate option of preparative treatment before allo-HCT for patients with AML in complete remission or MDS. Limitations of the meta-analysis included the small number of included clinical trials, significant heterogeneity between included studies for some outcomes, and lack of blinding in some studies.

Randomized Controlled Trials

No published randomized trials have compared RIC plus allo-HCT with conventional chemotherapy alone in patients with MDS and AML for whom MAC chemotherapy and allo-HCT are contraindicated.

Three RCTs, all of which are included in the systematic review by Song et al. (2021), (11) have compared RIC and myeloablative regimens before allo-HCT in patients with MDS. (12-14) The RCTs are heterogeneous in patient characteristics and conditioning regimens and their findings vary based on these differences. In a long-term follow-up of one of the RCTs (13), Scott et al. (2021) found that, at 4 years, transplant-related mortality was significantly increased with MAC as compared to RIC (25.1% vs. 9.9%; $p<.001$) and those who received RIC had a significantly increased relapse risk (HR, 4.06; 95% CI, 2.59 to 6.35; $p<.001$). (15) Among those who relapsed after HCT, postrelapse survival was similar between groups at 3 years (24% for MAC vs. 26% for RIC). Patients administered MAC had superior OS (HR, 1.54; 95% CI, 1.07 to 2.2; $p=.03$).

Overall, findings from these RCTs appear consistent with the American Society for Blood and Marrow Transplantation's (2009) systematic review (previously described), which assessed the evidence supporting reduced-intensity and myeloablative conditioning regimens and drew the following conclusions: "There are insufficient data to make a recommendation for an optimal conditioning regimen intensity. A range of dose intensities is currently being investigated, and the optimal approach will likely depend on disease and patient characteristics, such as age and comorbidities." (9) Other reviews (2010 to 2012) have also drawn conclusions similar to those of the American Society for Blood and Marrow Transplantation. (16-21) Given the absence of curative therapies for these patients, RIC allo-HCT may be considered as a risk-adapted treatment strategy for patients with MDS who could benefit from allo-HCT but who are at high risk of MAC regimen intolerance.

Noncomparative and Observational Studies

Additional nonrandomized evidence includes uncontrolled studies and prospective and retrospective cohort studies. Evidence from a number of largely heterogeneous, uncontrolled studies of RIC with allo-HCT has shown long-term remission (i.e., >4 years) can be achieved, often with reduced treatment-related morbidity and mortality, in patients with MDS or AML

who otherwise would not be candidates for MAC regimens. (8, 22-32) These prospective and retrospective studies included cohorts of 16 to 215 patients similar to those in the MAC allo-HCT studies. The most common conditioning regimens used were fludarabine-based, with cyclosporine A and tacrolimus used for GVHD prophylaxis. The reported incidence of grades II to IV GVHD was 9% to 63%, with a relapse risk of 6% to 61%. Rates of OS ranged between 44% at 1 year and 46% at 5 years (median follow-up range, 14 months to >4 years).

In general, nonrandomized studies of RIC compared to MAC showed a low rate of engraftment failure and low non-relapse mortality with RIC, but a higher relapse rate than with MAC allo-HCT. Zeng et al. (2014) conducted a systematic review and meta-analysis comparing outcomes for patients who had MDS or AML treated with HCT plus reduced-intensity or myeloablative conditioning. (33) Reviewers included 8 studies (2 prospective, 8 retrospective), with a total of 6464 AML or MDS patients. Of these, 171 received RIC and 4893 received MAC. Overall, RIC treated patients were older and more likely to have multiple comorbidities. In the pooled analysis, OS, RFS, and nonrelapse mortality did not differ significantly between patients receiving reduced-intensity and myeloablative conditioning. Relapse incidence was significantly lower in the MAC arm (odds ratio [OR] for RIC vs. MAC, 1.41; 95% CI, 1.24 to 1.59; $p < .001$).

Aoki et al. (2015) compared RIC with MAC in a retrospective cohort of 448 patients (age range, 50 to 69 years) with advanced MDS (refractory anemia with excess blasts or refractory anemia in transformation). (34) Of the total, 197 (44%) and 251 (56%) received myeloablative or RIC, respectively. The groups differed at baseline: patients who received RIC were significantly more likely to be 60 to 69 years old (vs. 50 to 59 years; 47% for RIC vs. 47% for MAC; $p = .001$), and less likely to receive an unrelated donor transplant (54% vs. 70%; $p = .001$). Three-year OS rates did not differ between groups (44.1% for RIC vs. 42.7% for MAC; $p = .330$). Although patients treated with RIC had a significantly lower 3-year cumulative incidence of nonrelapse mortality (25.6% vs. 37.9%; $p = .002$), they had a significantly higher 3-year incidence of relapse than patients treated with MAC (29.9% vs. 22.8%; $p = .029$).

Kim et al. (2012) published a phase 3 randomized trial (N=83 patients) comparing toxicity rates for 2 conditioning regimens (reduced CY, fludarabine, and anti-thymocyte globulin; standard CY anti-thymocyte globulin). (35) Four patients had MDS, and the remaining patients had severe aplastic anemia. Overall, the incidence of reported toxicities was lower in patients receiving the RIC regimen (23% vs. 55%; $p = .003$). Subgroup analyses showed no differences in the overall results based on differential diagnosis.

Outcomes After Allogeneic Hematopoietic Cell Transplantation in Mixed Myelodysplastic Syndrome Populations

Noncomparative and Observational Studies

A number of studies, primarily retrospective, continue to report outcomes from allo-HCT for MDS in a variety of patient populations and to evaluate the impact of specific patient, conditioning, and donor characteristics on outcomes; representative studies are summarized in Table 1.

Table 1. Case Series of Hematopoietic Cell Transplantation Treatment for Myelodysplastic Syndrome

Study	Patient Population	Type of HCT	Summary of Outcomes
Basquiera et al. (2015) (36)	52 pediatric patients with MDS	• Allo-HCT (59% with related donors) • Stem cell source: ○ Bone marrow, 63% ○ Peripheral blood, 26% ○ Umbilical cord blood, 11%	• 5-y DFS=50% • 5-y OS=55%
Boehm et al. (2014) (37)	60 adults with MDS or secondary AML	• Allo-HCT • MAC in 36 patients; RIC in 24 patients	10-y OS=46%
Damaj et al. (2014) (38)	128 adults with MDS: 40 received AZA before HCT and 88 who received BSC	RIC allo-HCT	• 3-y OS=53% in AZA group vs 53% in BSC group (p=.69) • 3-y RFS=37% in AZA group vs 42% in BSC group (p=.78) • 3-y NRM=20% in AZA group vs 23% in BSC group (p=.74)
Di Stasi et al. (2014) (39)	227 patients with MDS or AML	• Allo-HCT • Donor source: ○ Matched-related, 38% ○ Matched-unrelated, 48% ○ Haploidentical, 14%	3-y PFS for patients in remission: • 57% for matched-related • 45% for matched-unrelated • 41% for haploidentical (p=.417)
Onida et al. (2014) (40)	• 523 patients with MDS • IPSS cytogenetic risk group: ○ Good risk: 53.5% ○ Intermediate risk: 24.5% ○ Poor risk: 22%	• Allo-HCT • RIC in 12%	5-y OS based on IPSS cytogenetic risk group: • Good risk: 48% • Intermediate risk: 45% • Poor risk: 30%

Oran et al. (2014) (41)	256 patients with MDS - Pretreatment: - No cytoreductive Chemotherapy: 30.5% - Chemotherapy: 15.6% - HMA: 47.7% - Chemotherapy + HMA: 6.2%	- Allo-HCT - RIC in 36.7%	3-y EFS based on cytoreductive therapy: - No cytoreductive chemotherapy: 44.2% - Chemotherapy: 30.6% - HMA: 34.2% - Chemotherapy + HMA: 32.8% (p=.50)
Yoshimi et al. (2014) (42)	17 children with secondary MDS or AML after childhood aplastic anemia	Allo-HCT	5-y OS and EFS=41%
Basquiera et al. (2016) (43)	84 adults with MDS - Cytogenic risk group: - Standard: 65.5% - Adverse: 12.6% - Unknown: 21.9%	- Allo-HCT - RIC in 31.1%	OS: - Median: 23.5 mo (95% CI, 1.7 to 45.3 mo) 1-y=61% (95% CI, 50% to 70%) 4-y=38% (95% CI, 27% to 49%) PFS: - Median: 19.9 mo (95% CI, 9 to 31 mo) 1-y=57% (95% CI, 46% to 67%) 4-y=37% (95% CI, 26% to 48%)
Symeonidis et al. (2015) (44)	513 adults with CMML - Pretreatment: - No prior DMT: 28% - DMT: 72%	- Allo-HCT - RIC in 41.6%	1-y NRM=31% 4-y NRM=41% 4-y RFS=27% 4-y OS=33%
Pohlen et al. (2016) (45)	187 patients with refractory AML (87%) or high- risk MDS (13%)	- Allo-HCT - RIC in 52% - Unrelated donors in 73% - Stem cell source: - Bone marrow, 6%	3-y RFS=32% (95% CI, 25% to 39%) 3-y OS=35% (95% CI, 27% to 42%)

		○ Peripheral blood, 94%	
Heidenreich et al. (2017) (46)	• 313 adults with MDS and secondary AML, age ≥ 70 - Cytogenic risk group: ○ Good: 51% ○ Intermediate: 22% ○ Poor/very poor: 11%	• Allo-HCT • RIC or non-MAC in 83% • Unrelated donors in 75% • Stem cell source: ○ Bone marrow, 6% ○ Peripheral blood, 94%	• 1-y NRM=32% • 3-y relapse=28% • 3-y OS=34%
Robin et al. (2022) (47)	• 1114 adults with CMML age 18 to 70 years • CMML Prognosis Scoring System risk: ○ Low: 20% ○ Intermediate-1: 31% ○ Intermediate-2: 40% ○ High: 9% • Underwent allo-HCT: 43% • Transformed to AML prior to allo-HCT: 10%	• MAC or RIC allo-HCT; details of intensity and donor source not reported	• 5-y OS: ○ Lower-risk disease: 20% with allo-HCT vs. 42% without allo-HCT ($p<0.001$) ○ Higher-risk disease: 27% with allo-HCT vs. 15% without allo-HCT ($p=0.13$) • Multivariate analyses of risk of death within 2 years and after 2 years: ○ Lower-risk disease: Increased risk of death within 2 years with allo-HCT (HR=3.19); no difference in long-term survival after 2 years (HR=0.98)

			○ Higher-risk disease: Increased risk of death within 2 years with allo-HCT (HR=1.46); no difference in long-term survival after 2 years (HR=0.60) • Conditioning regimen and donor type were not associated with post-transplant survival (data not reported)
Sebert et al. (2025) (48)	• 77 adults with MDS (including CMML) age 18 to 69 years • Low to intermediate-1 risk per IPSS classification • And at least 1 additional risk factor: ○ Intermediate or higher IPSS risk ○ Thrombocytopenia <20 G/L ○ Neutropenia < 0.5 G/L ○ Failure of 2 lines of therapy	• Allo-HCT (n=72 with HLA-matched donors)	3-y OS: • 58.9% in HLA-matched donor arm vs 56.8% in non-donor arm (HR=0.75; p=.53) • Recruitment stopped for futility

allo: allogeneic; AML: acute myelogenous leukemia; AZA: azacitidine; BSC: best supportive care; chemo: chemotherapy; CI: confidence interval; CMML: chronic myelomonocytic leukemia; DFS: disease-free survival; DMT: disease-modifying therapy; EFS: event-free survival; HLA: human leukocyte antigen; HMA: hypomethylating agents; HCT: hematopoietic cell transplantation; HR: hazard ratio; IPSS: International Prognostic Scoring System; MAC: myeloablative conditioning; MDS: myelodysplastic syndrome; mo: month; NRM: non-relapse mortality; OS: overall survival; PFS: progression-free survival; RFS: relapse-free survival; RIC: reduced-intensity conditioning; y: year.

Section Summary: Myelodysplastic Syndrome

Primarily uncontrolled, observational studies of HCT for MDS have reported a relatively large range of OS and progression-free survival values, which reflect the heterogeneity in patient populations, conditioning regimens, and other factors. Reported estimates for 3- to 5-year OS of 40% to 50% are typical. Evidence from randomized and nonrandomized comparisons has suggested that RIC may be used as a risk-adapted strategy in high-risk patients who are older and with more comorbidities without significantly worsening OS. RIC appears to be associated with lower rates of nonrelapse mortality but higher cancer relapse than MAC HCT.

Myeloproliferative Neoplasms

Clinical Context and Therapy Purpose

The purpose of MAC and RIC allo-HCT in individuals who have myeloproliferative neoplasms is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The following PICO was used to select literature to inform this policy.

Populations

The relevant population of interest is individuals who have myeloproliferative neoplasms.

Interventions

The therapies being considered are MAC or RIC allo-HCT.

Comparators

The following therapies are currently being used: standard of care.

Outcomes

The general outcomes of interest are mortality and morbidity. Beneficial outcomes are an improvement in OS and disease-specific survival. Harmful outcomes are treatment-related morbidity and mortality. Follow-up over months to years is of interest for relevant outcomes.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs.
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

Data on therapy for myeloproliferative neoplasms are sparse. (29, 49, 50) As outlined in this medical policy, with the exception of MAC chemotherapy and allo-HCT, no therapy has yet proven to be curative or to prolong survival of patients with myeloproliferative neoplasms.

Systematic Reviews

Bewersdorf et al. (2021) assessed the available evidence on the efficacy and safety of allo-HCT in patients with myelofibrosis in a systematic review involving 43 studies (N=8739). (51) The analysis included 38 retrospective, 1 prospective, and 4 phase II clinical trials. Conditioning regimens used were variable with only 3 and 14 studies using exclusively MAC or RIC regimens, respectively. Additionally, donor sources and pre-transplantation treatment histories differed considerably among studies. The co-primary outcome was 1-, 2-, and 5-year OS. Rates of nonrelapse mortality, RFS or progression-free survival (PFS), and safety were also evaluated. Regarding survival, 1-year, 2-year, and 5-year OS rates were 66.7% (95% CI, 63.5% to 69.8%), 64.4% (95% CI, 57.6% to 70.6%), and 55% (95% CI, 51.8% to 58.3%), respectively. Nonrelapse mortality rates for the same time periods were 25.9% (95% CI, 23.3% to 28.7%), 29.7% (95% CI, 24.5% to 35.4%), and 30.5% (95% CI, 25.9% to 35.5%). Rates of 1-, 2- and 5-year RFS were 65.3% (95% CI, 56.5% to 73.1%), 56.2% (95% CI, 41.6% to 69.8%), and 53.6% (95% CI, 39.9% to 66.9%), respectively. PFS rates were 56.9% (95% CI, 41.4% to 71.2%), 50.6% (95% CI, 39.7% to 61.4%), and 43.5% (95% CI, 31.9% to 55.8%) for these same time periods. Acute GVHD was reported in 44% of patients, with chronic GVHD occurring in 46.5% of patients. The combined rate of graft failure was 10.6% (95% CI, 8.9% to 12.5%). Overall, the quality of the evidence was limited by the absence of RCTs and the retrospective design of most studies. Additionally, patient and transplant characteristics were variable among the included studies leading to moderate to substantial heterogeneity in the analyses.

Noncomparative and Observational Studies

The largest study identified evaluating allo-HCT for primary myelofibrosis comes from a 2010 analysis of the outcomes for 289 patients treated between 1989 and 2002, from the database of the Center for International Bone Marrow Transplant Research. (52) Median age was 47 years (range, 18 to 73 years). Donors were human leukocyte antigen (HLA)-identical siblings in 162 patients, unrelated individuals in 101 patients, and HLA nonidentical family members in 26 patients. Patients were treated with a variety of conditioning regimens and GVHD prophylaxis regimens. Splenectomy was performed in 65 patients before transplantation. The 100-day treatment-related mortality was 18% for HLA-identical sibling transplants, 35% for unrelated transplants, and 19% for transplants from alternative-related donors. Corresponding 5-year OS rates were 37%, 30%, and 40%, respectively. Disease-free survival (DFS) rates were 33%, 27%, and 22%, respectively. Rates of DFS for patients receiving reduced-intensity conditioning allo-HCT were comparable: 39% for HLA-identical sibling donors and 17% for unrelated donors at 3 years. In this large retrospective series, allogeneic transplantation for myelofibrosis resulted in long-term RFS in about one-third of patients.

The significant toxicity of MAC plus allo-HCT in myeloproliferative neoplasms has led to study of RIC regimens for these diseases. Data from a direct, prospective comparison of outcomes of MAC allo-HCT versus RIC allo-HCT in myeloproliferative neoplasms are not available, but single-arm series and nonrandomized comparative studies have reported outcomes after RIC allo-HCT. One 2008 series included 27 patients (mean age, 59 years) with myeloproliferative neoplasms who underwent allo-HCT using a RIC regimen of low-dose (2 gray) total body irradiation alone with or without fludarabine. (27) At a median follow-up of 47 months, 3-year RFS was 37%, 3-year OS was 43%, and 3-year nonrelapse mortality was 32%.

A 2009 retrospective study analyzed the impact of conditioning intensity on outcomes for allo-HCT in patients with myelofibrosis. (53) This multicenter trial included 46 consecutive patients treated at 3 Canadian and 4 European transplant centers between 1998 and 2005. Twenty-three patients (median age, 47 years; range, 31 to 60 years) underwent MAC and 23 patients (median age, 54 years; range, 38 to 74 years) underwent RIC. The majority in both groups (85%) were deemed intermediate- or high-risk. At a median follow-up of 50 months (range, 20 to 89 months), there was a trend for a better PFS rate at 3 years in RIC patients than in MAC patients (58% [range, 23% to 62%] vs. 43% [range, 35% to 76%], respectively; $p=.11$); there was a similar trend in the 3-year OS rate (68% [range, 45% to 84%] vs. 48% [range, 27% to 66%], respectively; $p=.08$). Nonrelapse mortality rates at 3 years trended higher in MAC cases (48%; range, 31% to 74%) than in RIC cases (27%; range, 14% to 55%; $p=.08$). The results of this study suggested that both types of conditioning regimens have curative potential in patients with myelofibrosis. Despite the RIC patients being significantly older, with longer disease duration and poorer performance status than those who received conventional conditioning, the groups had similar outcomes, supporting the use of RIC allo-HCT in this population.

Section Summary: Myeloproliferative Neoplasms

Observational studies of HCT for myeloproliferative neoplasms have reported a range of 3- to 5-year OS rates from 35% to 50% and suggested that HCT may be associated with improved survival in patients with intermediate-2 and high-risk disease. Primarily, retrospective studies have compared the RIC and MAC regimens. While these nonrandomized comparisons have suggested that RIC may be used in patients who are older and who have poorer performance status without significantly worsening OS, randomized trials are needed to provide greater certainty in the efficacy of the conditioning regimens.

Summary of Evidence

For individuals who have myelodysplastic syndrome (MDS) who receive myeloablative conditioning (MAC) or reduced-intensity conditioning (RIC) allogeneic hematopoietic cell transplantation (allo-HCT), the evidence includes systemic reviews, randomized controlled trials (RCTs), and numerous case series, which are often heterogeneous in terms of diseases included. Relevant outcomes are overall survival (OS), disease-specific survival, and treatment-related mortality and morbidity. Primarily uncontrolled, observational studies of hematopoietic cell transplantation (HCT) for MDS have reported a relatively large range of overall and progression free survival (PFS) rates, which reflect the heterogeneity in patient populations, conditioning regimens, and other factors. Reported estimates for 3- to 5-year OS of 40% to 50% are typical. Evidence from randomized and nonrandomized comparisons has suggested that RIC may be used as a risk-adapted strategy in high-risk patients who are older and with more comorbidities without significantly worsening OS. RIC appears to be associated with lower rates of nonrelapse mortality but higher cancer relapse than MAC HCT. At present, HCT is the only potentially curative treatment option for patients with MDS. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have myeloproliferative neoplasms who receive MAC or RIC allo-HCT, the evidence includes a systematic review and retrospective observational series. Relevant outcomes are OS, disease-specific survival, and treatment-related mortality and morbidity. Evidence has suggested that RIC may be used as a risk-adapted strategy in high-risk patients who are older and have more comorbidities without significantly worsening OS. RIC appears to be associated with lower rates of nonrelapse mortality but higher cancer relapse than myeloablative HCT. At present, HCT is the only potentially curative treatment option for patients with myeloproliferative neoplasms. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

Practice Guidelines and Position Statements

American Society of Transplantation and Cellular Therapy

In 2020, the American Society of Transplantation and Cellular Therapy (ASTCT) (formerly The American Society for Blood and Marrow Transplantation) published updated guidelines on indications for HCT and immune effector cell therapy based on the recommendations of a multiple-stakeholder task force. (54) Table 2 summarizes categorizations for allo-HCT in adults.

Table 2. Recommendations for the Use of Hematopoietic Cell Transplantation to Treat Myelodysplastic Syndromes, Myelofibrosis, and Myeloproliferative Neoplasms

Indication	Recommendation
Myelodysplastic Syndromes	
Low/intermediate-1 risk	Standard of care, clinical evidence available (large clinical trials and observational studies are not available; however, sufficiently large cohort studies have shown efficacy with “acceptable risk of morbidity and mortality”)
Intermediate-2/high-risk	Standard of care (“well defined and generally supported by evidence in the form of high-quality clinical trials and/or observational studies”)
Myelofibrosis and Myeloproliferative Neoplasms	
Primary, low risk	Standard of care (“well defined and generally supported by evidence in the form of high-quality clinical trials and/or observational studies”)
Primary, intermediate/high risk	Standard of care (“well defined and generally supported by evidence in the form of high-quality clinical trials and/or observational studies”)
Secondary	Standard of care (“well defined and generally supported by evidence in the form of high-quality clinical trials and/or observational studies”)
Hypereosinophilic syndromes, refractory	Standard of care, rare indication (clinical trials and observational studies are not feasible due to low incidence; small cohorts have shown efficacy with “acceptable risk of morbidity and mortality”)

In 2022, the ASTCT published guidelines on HCT in the treatment of acute myeloid leukemia (AML) and MDS in pediatric patients. (55) In 2023, the ASTCT published practice recommendations for HCT in the management of myelodysplastic syndromes in adults. (56) A standardized system for grading the levels of evidence was applied (as recommended by the ASTCT Steering Committee for evidence-based reviews). Table 3 summarizes allo-HCT specific recommendations by ASTCT for both pediatric and adult patients.

Table 3. Recommendations for the Use of Allogeneic Hematopoietic Cell Transplantation to Treat Myelodysplastic Syndromes

Indication/Consideration	Recommendation	Grade of Recommendation
<i>Adult patients</i>		
Should allogeneic HCT routinely be offered early for advanced (int-2/high) de novo MDS?	Yes	A
Should allogeneic HCT routinely be offered early for lower risk (low/int-1) de novo MDS?	No	B
<i>Pediatric patients</i>		
Should children with favorable risk cytogenetic and molecular lesions (<i>CBF</i> , <i>NPM1</i> , <i>CEBPA bZip</i>) undergo HCT in CR1, even if MRD ⁺ at first end-of-induction?	No	A
Should children with <i>FLT3-ITD</i> mutations undergo HCT in CR1?	Yes	B
Should children with high-risk cytomolecular abnormalities undergo HCT in CR1?	Yes	B, C, or D (variant-dependent)
Should children who are MRD ⁺ by flow cytometry at end-of-induction and no other risk-stratifying lesion be considered for HCT in CR1?	Yes	A
Should children with primary induction failure or refractory disease after 2-3 cycles of chemotherapy be considered for allogeneic HCT, even if not in CR1?	Yes	B
Should children who relapse be offered HCT following attempts to obtain CR2?	Yes	B
For children with inherited bone marrow failure syndromes who develop MDS, should reduced-toxicity or reduced-intensity conditioning be favored?	Yes	C
Should myeloablative conditioning be used for patients undergoing first or second HCT assuming a suitable donor and that comorbidities do not warrant a reduced-intensity approach?	Yes	B

CR1: first complete remission; CR2: second complete remission; HCT: hematopoietic cell transplantation; MDS: myelodysplastic syndrome; MRD: measurable residual disease; RCT: randomized controlled trial. Grades of Recommendation: A = at least 1 meta-analysis, systematic review, or RCT rated as 1++ and directly applicable to the target population or a systematic review of RCTs or a body of evidence consisting principally of studies rated as 1+ directly applicable to the target population and demonstrating overall consistency of results; B = a body of evidence including studies rated as 2++ directly applicable to the target population and demonstrating overall consistency of results or extrapolated evidence from studies rated as 1++ or 1+; C = a body of evidence including studies rated as 2+ directly applicable to the target population and demonstrating overall consistency of results or extrapolated evidence from studies rated as 2++; D = evidence level 3 or 4 or extrapolated evidence from studies rated as 2+.

Relevant levels of evidence: 1++ = high quality meta-analyses, systematic reviews of RCTs, or RCTs with a very low risk of bias; 1+ = well conducted meta-analyses, systematic reviews of RCTs, or RCTs with a low risk of bias; 2++ = high quality systematic reviews of case-control or cohort studies or high quality case-control or cohort studies with a very low risk of confounding, bias, or chance and a high probability that the relationship is causal; 2+ = well conducted case-control or cohort studies with a low risk of confounding, bias, or chance and a moderate probability that the relationship is causal; 3 = non-analytic studies, e.g., case reports, case series; 4 = expert opinion.

National Comprehensive Cancer Network

Current National Comprehensive Cancer Network clinical guidelines for myelodysplastic syndromes (v.3.2026) make the following general recommendation about allo-HCT (57):

“For patients who are transplant candidates, an HLA [human leukocyte antigen]-matched sibling, or HLA-matched unrelated donor can be considered. Results with HLA-matched unrelated donors have improved to levels comparable to those obtained with HLA-matched siblings. With the increasing use of cord blood or HLA-haploidentical related donors, HCT has become a viable option for many patients. High-dose conditioning is typically used for younger patients, whereas RIC [reduced-intensity conditioning] for HCT is generally the strategy in older individuals.”

Specific National Comprehensive Cancer Network recommendations for HCT for treatment of myelodysplastic syndromes are outlined in Table 4.

Table 4. Guidelines for Allogeneic Hematopoietic Cell Transplantation for Myelodysplastic Syndromes

Prognostic Category	Recommendations for Allo-HCT
IPSS low/intermediate-1 <i>OR</i> IPSS-R very low, low, intermediate <i>OR</i> WPSS very low, low, intermediate	- Consider allo-HCT for select patients who have clinically relevant thrombocytopenia or neutropenia, with disease progression or no response after azacitidine/decitabine or immunosuppressive therapy - Consider allo-HCT for patients who have symptomatic anemia with no 5q deletion, with serum erythropoietin level >500 mU/mL or

	lower serum erythropoietin level with inadequate response to erythropoietin stimulating agents and/or lenalidomide, with poor probability of or inadequate response/intolerance to immunosuppressive therapy, and no response or intolerance to azacitidine/decitabine or immunosuppressive therapy Consider allo-HCT for patients who have symptomatic anemia with del(5q), with inadequate response/intolerance to lenalidomide and/or erythropoietin stimulating agents, and no response or intolerance to azacitidine/decitabine or immunosuppressive therapy
IPSS intermediate-2, high <i>OR</i> IPSS-R intermediate, high, very high <i>OR</i> WPSS high, very high	Recommend allo-HCT if a high-intensity therapy candidate and transplant candidate and donor stem cell source is available

allo: allogeneic; HCT: hematopoietic cell transplantation; IPSS: International Prognostic Scoring System; WPSS: WHO Classification-based Prognostic Scoring System.

Table 5 summarizes the NCCN recommendations (v.1.2026) on the use of allo-HCT for the treatment of myeloproliferative neoplasms. (58) The guidelines note that selection of allo-HCT should be based on age, performance status, major comorbid conditions, psychosocial status, patient preference, and availability of caregiver.

Table 5. Guidelines for Allogeneic Hematopoietic Cell Transplantation for Myeloproliferative Neoplasms

Prognostic Category	Recommendations for Allo-HCT
Lower-risk myelofibrosis MIPSS-70: ≤ 3 MIPSS-70+: Version 2.0 ≤ 3 DIPSS-Plus: ≤ 1 DIPSS: ≤ 2 MYSEC-PM: < 14	- In symptomatic patients with disease progression despite treatment with ruxolitinib, peginterferon alfa-2a, or hydroxyurea (if cytoreduction would be symptomatically beneficial), pacritinib, or momelotinib, consider allo-HCT immediately or bridging therapy to decrease marrow blasts to an acceptable level prior to transplant - Evaluation for allo-HCT is recommended for patients with low platelet counts or complex cytogenetics
Higher-risk myelofibrosis MIPSS-70: ≥ 4 MIPSS-70+: Version 2.0 ≥ 4 DIPSS-Plus: > 1 DIPSS: > 2 MYSEC-PM: ≥ 14	- Consider allo-HCT immediately or bridging therapy can be used to decrease marrow blasts to an acceptable level prior to transplant - Evaluation for allo-HCT is recommended for all patients

Disease progression to advanced-stage/AML	• Bridging therapy followed by allo-HCT; bridging therapy options include: hypomethylating agents ± JAK inhibitors or venetoclax, or intensive induction chemotherapy, followed by allo-HCT
--	---

allo: allogeneic; AML: acute myeloid leukemia; DIPSS: Dynamic International Prognostic Scoring System; HCT: hematopoietic cell transplantation; MIPSS: Mutation-Enhanced International Prognostic Scoring System; MYSEC-PM: Myelofibrosis Secondary to PV [polycythemia vera] and ET [essential thrombocythemia]-Prognostic Model; JAK: Janus kinase.

Specific NCCN recommendations for both myelodysplastic syndromes and myeloproliferative neoplasms can be found at <<https://www.nccn.org>>.

Medicare National Coverage

There is a national coverage determination for stem cell transplantation (110.23; formerly 110.81), (59) portions of which are highlighted below:

Nationally Covered Indications:

- Allogeneic Hematopoietic Stem Cell Transplantation (HSCT)
 - "...Treatment of leukemia, leukemia in remission, or aplastic anemia when it is reasonable and necessary,
 - ...Treatment of severe combined immunodeficiency disease (SCID) and for the treatment of Wiskott-Aldrich syndrome,
 - ...Treatment of Myelodysplastic Syndromes (MDS) pursuant to Coverage with Evidence Development (CED) in the context of a Medicare-approved, prospective clinical study."

Medicare payment for these beneficiaries will be restricted to patients enrolled in an approved clinical study.

- "Effective ... January 27, 2016, allogeneic HSCT for multiple myeloma is covered by Medicare only for beneficiaries with Durie-Salmon Stage II or III multiple myeloma, or International Staging System (ISS) Stage II or Stage III multiple myeloma and participating in an approved prospective clinical study that meets the criteria below. There must be appropriate statistical techniques to control for selection bias and confounding by age, duration of diagnosis, disease classification, International Myeloma Working Group (IMWG) classification, ISS stage, comorbid conditions, type of preparative/conditioning regimen, graft versus host disease (GVHD) prophylaxis, donor type and cell source....
- Effective ... January 27, 2016, allogeneic HSCT for myelofibrosis (MF) is covered by Medicare only for beneficiaries with Dynamic International Prognostic Scoring System (DIPSSplus) intermediate-2 or High primary or secondary MF and participating in an approved prospective clinical study. All Medicare-approved studies must use appropriate statistical techniques in the analysis to control for selection bias and potential confounding by age, duration of diagnosis, disease classification, DIPSSplus score, comorbid conditions, type of preparative/conditioning regimen, graft vs. host disease (GVHD) prophylaxis, donor type and cell source....

- Effective ... January 27, 2016, allogeneic HSCT for sickle cell disease (SCD) is covered by Medicare only for beneficiaries with severe, symptomatic SCD who participate in an approved prospective clinical study....”

Ongoing and Unpublished Clinical Trials

Some currently ongoing trials that might influence this policy are listed in Table 6.

Table 6. Summary of Key Trials

NCT Number	Trial Name	Planned Enrollment	Completion Date
NCT05367583	Cohort Study Assessing the Treatment Strategy for High-Risk Myelodysplastic Syndromes in Patients Under 70 (COMYRE)	107	Oct 2024

NCT: national clinical trial.

Coding

Procedure codes on Medical Policy documents are included **only** as a general reference tool for each policy. **They may not be all-inclusive.**

The presence or absence of procedure, service, supply, or device codes in a Medical Policy document has no relevance for determination of benefit coverage for members or reimbursement for providers. **Only the written coverage position in a Medical Policy should be used for such determinations.**

Benefit coverage determinations based on written Medical Policy coverage positions must include review of the member’s benefit contract or Summary Plan Description (SPD) for defined coverage vs. non-coverage, benefit exclusions, and benefit limitations such as dollar or duration caps.

CPT Codes	36511, 38204, 38205, 38206, 38207, 38208, 38209, 38210, 38211, 38212, 38213, 38214, 38215, 38220, 38221, 38222, 38230, 38232, 38240, 38241, 38242, 38243, 81265, 81266, 81267, 81268, 81370, 81371, 81372, 81373, 81374, 81375, 81376, 81377, 81378, 81379, 81380, 81381, 81382, 81383, 86805, 86806, 86807, 86808, 86812, 86813, 86816, 86817, 86821, 86825, 86826, 86828, 86829, 86830, 86831, 86832, 86833, 86834, 86835, 86849, 86950, 86985, 88240, 88241
HCPCS Codes	S2140, S2142, S2150

*Current Procedural Terminology (CPT®) ©2025 American Medical Association: Chicago, IL.

References

1. Khoury JD, Solary E, Abla O, et al. The 5th edition of the World Health Organization Classification of Haematolymphoid Tumours: Myeloid and Histiocytic/Dendritic Neoplasms. Leukemia. Jul 2022; 36(7):1703-1719. PMID 35732831
2. Schanz J, Tüchler H, Solé F, et al. New comprehensive cytogenetic scoring system for primary myelodysplastic syndromes (MDS) and oligoblastic acute myeloid leukemia after

- MDS derived from an international database merge. *J Clin Oncol*. Mar 10 2012; 30(8):820-829. PMID 22331955
3. Verstovsek S, Mesa RA, Gotlib J, et al. A double-blind, placebo-controlled trial of ruxolitinib for myelofibrosis. *N Engl J Med*. Mar 01 2012; 366(9):799-807. PMID 22375971
 4. Cervantes F, Vannucchi AM, Kiladjan JJ, et al. Three-year efficacy, safety, and survival findings from COMFORT-II, a phase 3 study comparing ruxolitinib with best available therapy for myelofibrosis. *Blood*. Dec 12 2013; 122(25):4047-4053. PMID 24174625
 5. Harrison C, Kiladjan JJ, Al-Ali HK, et al. JAK inhibition with ruxolitinib versus best available therapy for myelofibrosis. *N Engl J Med*. Mar 01 2012; 366(9):787-798. PMID 22375970
 6. Food and Drug Administration. FDA approves fedratinib for myelofibrosis. August 2019. Available at <<https://www.fda.gov>> (accessed November 25, 2025).
 7. Kasner MT, Luger SM. Update on the therapy for myelodysplastic syndrome. *Am J Hematol*. Mar 2009; 84(3):177-186. PMID 19195035
 8. Kindwall-Keller T, Isola LM. The evolution of hematopoietic SCT in myelodysplastic syndrome. *Bone Marrow Transplant*. Apr 2009; 43(8):597-609. PMID 19252532
 9. Oliansky DM, Antin JH, Bennett JM, et al. The role of cytotoxic therapy with hematopoietic stem cell transplantation in the therapy of myelodysplastic syndromes: an evidence-based review. *Biol Blood Marrow Transplant*. Feb 2009; 15(2):137-172. PMID 19167676
 10. Koenecke C, Göhring G, de Wreede LC, et al. Impact of the revised International Prognostic Scoring System, cytogenetics and monosomal karyotype on outcome after allogeneic stem cell transplantation for myelodysplastic syndromes and secondary acute myeloid leukemia evolving from myelodysplastic syndromes: a retrospective multicenter study of the European Society of Blood and Marrow Transplantation. *Haematologica*. Mar 2015; 100(3):400-408. PMID 25552702
 11. Song Y, Yin Z, Ding J, et al. Reduced Intensity Conditioning Followed by Allogeneic Hematopoietic Stem Cell Transplantation Is a Good Choice for Acute Myeloid Leukemia and Myelodysplastic Syndrome: A Meta-Analysis of Randomized Controlled Trials. *Front Oncol*. 2021; 11:708727. PMID 34692485
 12. Beelen DW, Trensche R, Stelljes M, et al. Treosulfan or busulfan plus fludarabine as conditioning treatment before allogeneic haemopoietic stem cell transplantation for older patients with acute myeloid leukaemia or myelodysplastic syndrome (MC-FludT.14/L): a randomised, non-inferiority, phase 3 trial. *Lancet Haematol*. Jan 2020; 7(1):e28-e39. PMID 31606445
 13. Scott BL, Pasquini MC, Logan BR, et al. Myeloablative Versus Reduced-Intensity Hematopoietic Cell Transplantation for Acute Myeloid Leukemia and Myelodysplastic Syndromes. *J Clin Oncol*. Apr 10 2017; 35(11):1154-1161. PMID 28380315
 14. Kröger N, Iacobelli S, Franke GN, et al. Dose-Reduced Versus Standard Conditioning Followed by Allogeneic Stem-Cell Transplantation for Patients With Myelodysplastic Syndrome: A Prospective Randomized Phase III Study of the EBMT (RICMAC Trial). *J Clin Oncol*. Jul 01 2017; 35(19):2157-2164. PMID 28463633
 15. Scott BL, Pasquini MC, Fei M, et al. Myeloablative versus Reduced-Intensity Conditioning for Hematopoietic Cell Transplantation in Acute Myelogenous Leukemia and Myelodysplastic Syndromes-Long-Term Follow-Up of the BMT CTN 0901 Clinical Trial. *Transplant Cell Ther*. Jun 2021; 27(6):483.e1-483.e6. PMID 33775615

16. Akhtari M. When to treat myelodysplastic syndromes. *Oncology (Williston Park)*. May 2011; 25(6):480-486. PMID 21717901
17. Deeg HJ, Sandmaier BM. Who is fit for allogeneic transplantation? *Blood*. Dec 02 2010; 116(23):4762-4770. PMID 20702782
18. Giralt SA, Horowitz M, Weisdorf D, et al. Review of stem-cell transplantation for myelodysplastic syndromes in older patients in the context of the Decision Memo for Allogeneic Hematopoietic Stem Cell Transplantation for Myelodysplastic Syndrome emanating from the Centers for Medicare and Medicaid Services. *J Clin Oncol*. Feb 10 2011; 29(5):566-572. PMID 21220586
19. Deeg HJ, Bartenstein M. Allogeneic hematopoietic cell transplantation for myelodysplastic syndrome: current status. *Arch Immunol Ther Exp (Warsz)*. Feb 2012; 60(1):31-41. PMID 22143157
20. Garcia-Manero G. Myelodysplastic syndromes: 2012 update on diagnosis, risk-stratification, and management. *Am J Hematol*. Jul 2012; 87(7):692-701. PMID 22696212
21. Kröger N. Allogeneic stem cell transplantation for elderly patients with myelodysplastic syndrome. *Blood*. Jun 14 2012; 119(24):5632-5639. PMID 22504927
22. Barrett AJ, Savani BN. Allogeneic stem cell transplantation for myelodysplastic syndrome. *Semin Hematol*. Jan 2008; 45(1):49-59. PMID 18179969
23. Blaise D, Vey N, Faucher C, et al. Current status of reduced-intensity-conditioning allogeneic stem cell transplantation for acute myeloid leukemia. *Haematologica*. Apr 2007; 92(4):533-541. PMID 17488664
24. Deschler B, de Witte T, Mertelsmann R, et al. Treatment decision-making for older patients with high-risk myelodysplastic syndrome or acute myeloid leukemia: problems and approaches. *Haematologica*. Nov 2006; 91(11):1513-1522. PMID 17082009
25. Huisman C, Meijer E, Petersen EJ, et al. Hematopoietic stem cell transplantation after reduced intensity conditioning in acute myelogenous leukemia patients older than 40 years. *Biol Blood Marrow Transplant*. Feb 2008; 14(2):181-186. PMID 18215778
26. Kröger N, Bornhäuser M, Ehninger G, et al. Allogeneic stem cell transplantation after a fludarabine/busulfan-based reduced-intensity conditioning in patients with myelodysplastic syndrome or secondary acute myeloid leukemia. *Ann Hematol*. Jun 2003; 82(6):336-342. PMID 12728337
27. Laport GG, Sandmaier BM, Storer BE, et al. Reduced-intensity conditioning followed by allogeneic hematopoietic cell transplantation for adult patients with myelodysplastic syndrome and myeloproliferative disorders. *Biol Blood Marrow Transplant*. Feb 2008; 14(2):246-255. PMID 18215785
28. Martino R, Caballero MD, Pérez-Simón JA, et al. Evidence for a graft-versus-leukemia effect after allogeneic peripheral blood stem cell transplantation with reduced-intensity conditioning in acute myelogenous leukemia and myelodysplastic syndromes. *Blood*. Sep 15 2002; 100(6):2243-2245. PMID 12200391
29. Mesa RA. Navigating the evolving paradigms in the diagnosis and treatment of myeloproliferative disorders. *Hematology Am Soc Hematol Educ Program*. 2007:355-362. PMID 18024651
30. Tauro S, Craddock C, Peggs K, et al. Allogeneic stem-cell transplantation using a reduced-intensity conditioning regimen has the capacity to produce durable remissions and long-

- term disease-free survival in patients with high-risk acute myeloid leukemia and myelodysplasia. *J Clin Oncol.* Dec 20 2005; 23(36):9387-9393. PMID 16314618
31. Valcárcel D, Martino R. Reduced intensity conditioning for allogeneic hematopoietic stem cell transplantation in myelodysplastic syndromes and acute myelogenous leukemia. *Curr Opin Oncol.* Nov 2007; 19(6):660-666. PMID 17906468
 32. Valcárcel D, Martino R, Caballero D, et al. Sustained remissions of high-risk acute myeloid leukemia and myelodysplastic syndrome after reduced-intensity conditioning allogeneic hematopoietic transplantation: chronic graft-versus-host disease is the strongest factor improving survival. *J Clin Oncol.* Feb 01 2008; 26(4):577-584. PMID 18086801
 33. Zeng W, Huang L, Meng F, et al. Reduced-intensity and myeloablative conditioning allogeneic hematopoietic stem cell transplantation in patients with acute myeloid leukemia and myelodysplastic syndrome: a meta-analysis and systematic review. *Int J Clin Exp Med.* 2014; 7(11):4357-4368. PMID 25550955
 34. Aoki K, Ishikawa T, Ishiyama K, et al. Allogeneic haematopoietic cell transplantation with reduced-intensity conditioning for elderly patients with advanced myelodysplastic syndromes: a nationwide study. *Br J Haematol.* Feb 2015; 168(3):463-466. PMID 25228239
 35. Kim H, Lee JH, Joo YD, et al. A randomized comparison of cyclophosphamide vs. reduced dose cyclophosphamide plus fludarabine for allogeneic hematopoietic cell transplantation in patients with aplastic anemia and hypoplastic myelodysplastic syndrome. *Ann Hematol.* Sep 2012; 91(9):1459-1469. PMID 22526363
 36. Basquiera AL, Pizzi S, Correias AG, et al. Allogeneic hematopoietic stem cell transplantation in pediatric myelodysplastic syndromes: a multicenter experience from Argentina. *Pediatr Blood Cancer.* Jan 2015; 62(1):153-157. PMID 25264233
 37. Boehm A, Sperr WR, Kalhs P, et al. Long-term follow-up after allogeneic stem cell transplantation in patients with myelodysplastic syndromes or secondary acute myeloid leukemia: a single center experience. *Wien Klin Wochenschr.* Jan 2014; 126(1-2):23-29. PMID 24249320
 38. Damaj G, Mohty M, Robin M, et al. Upfront allogeneic stem cell transplantation after reduced-intensity/nonmyeloablative conditioning for patients with myelodysplastic syndrome: a study by the Société Française de Greffe de Moelle et de Thérapie Cellulaire. *Biol Blood Marrow Transplant.* Sep 2014; 20(9):1349-1355. PMID 24838178
 39. Di Stasi A, Milton DR, Poon LM, et al. Similar transplantation outcomes for acute myeloid leukemia and myelodysplastic syndrome patients with haploidentical versus 10/10 human leukocyte antigen-matched unrelated and related donors. *Biol Blood Marrow Transplant.* Dec 2014; 20(12):1975-1981. PMID 25263628
 40. Onida F, Brand R, van Biezen A, et al. Impact of the International Prognostic Scoring System cytogenetic risk groups on the outcome of patients with primary myelodysplastic syndromes undergoing allogeneic stem cell transplantation from human leukocyte antigen-identical siblings: a retrospective analysis of the European Society for Blood and Marrow Transplantation-Chronic Malignancies Working Party. *Haematologica.* Oct 2014; 99(10):1582-1590. PMID 25085359
 41. Oran B, Kongtim P, Popat U, et al. Cytogenetics, donor type, and use of hypomethylating agents in myelodysplastic syndrome with allogeneic stem cell transplantation. *Biol Blood Marrow Transplant.* Oct 2014; 20(10):1618-1625. PMID 24953017

42. Yoshimi A, Strahm B, Baumann I, et al. Hematopoietic stem cell transplantation in children and young adults with secondary myelodysplastic syndrome and acute myelogenous leukemia after aplastic anemia. *Biol Blood Marrow Transplant.* Mar 2014; 20(3):425-429. PMID 24316460
43. Basquiera AL, Rivas MM, Remaggi G, et al. Allogeneic hematopoietic stem cell transplantation in adults with myelodysplastic syndrome: Experience of the Argentinean Group of Bone Marrow Transplantation (GATMO). *Hematology.* Apr 2016; 21(3):162-169. PMID 26147089
44. Symeonidis A, van Biezen A, de Wreede L, et al. Achievement of complete remission predicts outcome of allogeneic haematopoietic stem cell transplantation in patients with chronic myelomonocytic leukaemia. A study of the Chronic Malignancies Working Party of the European Group for Blood and Marrow Transplantation. *Br J Haematol.* Oct 2015; 171(2):239-246. PMID 26212516
45. Pohlen M, Groth C, Sauer T, et al. Outcome of allogeneic stem cell transplantation for AML and myelodysplastic syndrome in elderly patients (≥ 60 years). *Bone Marrow Transplant.* Nov 2016; 51(11):1441-1448. PMID 27295269
46. Heidenreich S, Ziagkos D, de Wreede LC, et al. Allogeneic Stem Cell Transplantation for Patients Age ≥ 70 Years with Myelodysplastic Syndrome: A Retrospective Study of the MDS Subcommittee of the Chronic Malignancies Working Party of the EBMT. *Biol Blood Marrow Transplant.* Jan 2017; 23(1):44-52. PMID 27720995
47. Robin M, de Wreede LC, Padron E, et al. Role of allogeneic transplantation in chronic myelomonocytic leukemia: an international collaborative analysis. *Blood.* Sep 22 2022; 140(12):1408-1418. PMID 35667047
48. Sebert M, Thepot S, Cluzeau T, et al. Transplantation in lower risk MDS patients: a prospective phase 2 trial based on donor availability. *Blood Adv.* Jan 27 2026; 10(2):494-504. PMID 41061188
49. Tefferi A, Vainchenker W. Myeloproliferative neoplasms: molecular pathophysiology, essential clinical understanding, and treatment strategies. *J Clin Oncol.* Feb 10 2011; 29(5):573-582. PMID 21220604
50. McLornan DP, Mead AJ, Jackson G, et al. Allogeneic stem cell transplantation for myelofibrosis in 2012. *Br J Haematol.* May 2012; 157(4):413-425. PMID 22463701
51. Bewersdorf JP, Sheth AH, Vetsa S, et al. Outcomes of Allogeneic Hematopoietic Cell Transplantation in Patients With Myelofibrosis-A Systematic Review and Meta-Analysis. *Transplant Cell Ther.* Oct 2021; 27(10):873.e1-e873.e13. PMID 34052505
52. Ballen KK, Shrestha S, Sobocinski KA, et al. Outcome of transplantation for myelofibrosis. *Biol Blood Marrow Transplant.* Mar 2010; 16(3):358-367. PMID 19879949
53. Gupta V, Kröger N, Aschan J, et al. A retrospective comparison of conventional intensity conditioning and reduced-intensity conditioning for allogeneic hematopoietic cell transplantation in myelofibrosis. *Bone Marrow Transplant.* Sep 2009; 44(5):317-320. PMID 19234505
54. Kanate AS, Majhail NS, Savani BN, et al. Indications for Hematopoietic Cell Transplantation and Immune Effector Cell Therapy: Guidelines from the American Society for Transplantation and Cellular Therapy. *Biol Blood Marrow Transplant.* Jul 2020; 26(7):1247-1256. PMID 32165328

55. Tarlock K, Sulis ML, Chewning JH, et al. Hematopoietic Cell Transplantation in the Treatment of Pediatric Acute Myelogenous Leukemia and Myelodysplastic Syndromes: Guidelines from the American Society of Transplantation and Cellular Therapy. *Transplant Cell Ther.* Sep 2022; 28(9):530-545. PMID 35717004
56. DeFilipp Z, Ciurea SO, Cutler C, et al. Hematopoietic Cell Transplantation in the Management of Myelodysplastic Syndrome: An Evidence-Based Review from the American Society for Transplantation and Cellular Therapy Committee on Practice Guidelines. *Transplant Cell Ther.* February 2023; 29(2):71-81. PMID 36436780
57. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Myelodysplastic Syndromes, Version 3.2026. Available at <<https://www.nccn.org>> (accessed April 13, 2026).
58. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Myeloproliferative Neoplasms, Version 1.2026. Available at <<https://www.nccn.org>> (accessed April 13, 2026).
59. Centers for Medicare and Medicaid Services. National Coverage Determination (NCD) for Stem Cell Transplantation Formerly 110.8.1 (110.23). 2024. Available at <<https://www.cms.gov>> (accessed November 25, 2025).

Centers for Medicare and Medicaid Services (CMS)

The information contained in this section is for informational purposes only. HCSC makes no representation as to the accuracy of this information. It is not to be used for claims adjudication for HCSC Plans.

The Centers for Medicare and Medicaid Services (CMS) does have a national Medicare coverage position. Coverage may be subject to local carrier discretion.

A national coverage position for Medicare may have been changed since this medical policy document was written. See Medicare's National Coverage at <<https://www.cms.hhs.gov>>.

Policy History/Revision	
Date	Description of Change
05/15/2026	Document updated. Coverage unchanged. References 48 and 55 added; others updated.
12/15/2025	Document updated. The following changes were made to Coverage: 1) Moved information on myelodysplastic syndromes (MDS) and myeloproliferative neoplasms (MPN) categorization and risk for progression to the Policy Guidelines section; and 2) Removed statement on autologous hematopoietic cell transplantation. Added reference 57; others updated. Title changed from: Hematopoietic Cell Transplantation for Myelodysplastic Syndromes (MDS) and Myeloproliferative Neoplasms (MPN).
04/01/2025	Reviewed. No changes.

01/01/2025	Document updated with literature review. Coverage unchanged. Reference 56 added; some updated and others removed.
11/01/2023	Document updated with literature review. Coverage unchanged. References 1, 47, and 54 added; some updated and others removed.
05/15/2022	Document updated with literature review. Coverage unchanged. References 5, 10, 14, 48, and 53 added; some updated and others removed.
08/01/2021	Reviewed. No changes.
11/01/2020	Document updated with literature review. The following changes were made to the Coverage section: 1. The medically necessary statement for allogeneic hematopoietic cell transplantation (allo-HCT; HCT) to treat myelodysplastic syndromes (MDS) includes older age and Multiple comorbidities (e.g., HCT-comorbidity index [HCT-CI] score higher than 2) under any of the following indications. 2. NOTE 4 was added to define the HCT-CI. References 9-11 and 54 were added and others revised.
09/15/2019	Reviewed. No changes.
05/15/2018	Document updated with literature review. Coverage unchanged. References 38-39 and 49-50, 52 added.
06/01/2017	Reviewed. No changes.
10/15/2016	Document updated with literature review. The following was added to the allogeneic hematopoietic stem-cell transplantation medically necessary criteria for increasing number of blasts: "Refractory anemia with excess blasts; Refractory anemia with excess blasts in transformation; or Chronic myelomonocytic leukemia (CMML). The following NOTES and Tables were added to coverage: 1) International Prognostic Scoring System (IPSS) Variables and Outcome information for myelodysplastic syndromes (MDS); 2) Karotype definitions; and 3) Cytopenia definitions.
07/15/2015	Document updated with literature review. The following was added to the medically necessary criteria for myelodysplastic syndromes: "In patients with any of the following indications: International Prognostic Scoring System of intermediate-2 or high risk; Red blood cell transfusion dependence, neutropenia; Neutropenia; Thrombocytopenia; High-risk cytogenetics; or Increasing blast percentage." The following was added to the medically necessary criteria for myeloproliferative neoplasms: "When there are any of the following indications: Cytopenias; Transfusion dependence; Increasing blast percentage over 5%; or Age 30 to 65 years." The following coverage statement was added: "Allogeneic HSCT is considered experimental, investigational and/or unproven for myelodysplastic syndrome or for myeloproliferative neoplasm that does not meet the criteria listed above." Title changed from Stem-Cell Transplant for Myelodysplastic Syndromes (MDS) and Myeloproliferative Neoplasms (MPN).
06/01/2014	Document updated with literature review. The following was changed: 1) Expanded coverage to consider a) donor leukocyte infusion (DLI) and hematopoietic progenitor cell (HPC) boost as medically necessary for

	myelodysplastic syndromes and myeloproliferative neoplasms that has relapsed, to prevent relapse in the setting of a high-risk relapse, or to convert a patient from mixed to full donor chimerism; b) DLI and HPC boost are considered experimental, investigational and/or unproven following an AlloSCS treatment for MDS/MPN that was originally considered experimental, investigational and/or unproven for the treatment of MDS/MPN OR as a treatment prior to AlloSCS; and, 2) Expanded coverage to consider a) short tandem repeat (STR) markers medically necessary when used in pre- or post-stem-cell support testing of the donor and recipient DNA profiles as a way to assess the status of donor cell engraftment following AlloSCS for MDS/MPN; b) all other uses of STR markers MDS and MPN experimental, investigational and/or unproven, if not listed in the coverage section. Description and Rationale were significantly changed, including the 2008 World Health Organization Classification and International Prognostic Scoring System. Title changed from Stem-Cell Transplant for Myelodysplastic Syndromes and Myeloproliferative Diseases.
04/01/2010	New medical document originating from: SUR703.017, Peripheral/Bone Marrow Stem-Cell Transplantation (PSCT/BMT) for Non-Malignancies; SUR703.018, Peripheral/Bone Marrow Stem-Cell Transplantation (PSCT/BMT) for Malignancies; SUR703.022, Cord Blood as a Source of Stem-Cells (CBSC); SUR703.023, Donor Leukocyte Infusion (DLI); and SUR703.024, Tandem/Triple High-Dose Chemoradiotherapy with Stem-Cell Support for Malignancies. Stem-Cell transplant continues to be medically necessary when stated criteria are met. [NOTE: A link to the medical policies with the following titles can be found at the end of the medical policy SUR703.002, Stem-Cell Reinfusion or Transplantation Following Chemotherapy (General Donor and Recipient Information): • Peripheral/Bone Marrow Stem-Cell Transplantation (PSCT/BMT) for Non-Malignancies; • Peripheral/Bone Marrow Stem-Cell Transplantation (PSCT/BMT) for Malignancies; • Cord Blood as a Source of Stem-Cells; • Donor Leukocyte Infusion (DLI); and Tandem/Triple High-Dose Chemoradiotherapy with Stem-Cell Support for Malignancies.