

Policy Number	THE801.030
Policy Effective Date	05/01/2026

Nonpharmacologic Treatment of Rosacea

Table of Contents
Coverage
Policy Guidelines
Description
Rationale
Coding
References
Policy History

Related Policies (if applicable)
None

Disclaimer

Carefully check state regulations and/or the member contract.

Each benefit plan, summary plan description or contract defines which services are covered, which services are excluded, and which services are subject to dollar caps or other limitations, conditions or exclusions. Members and their providers have the responsibility for consulting the member's benefit plan, summary plan description or contract to determine if there are any exclusions or other benefit limitations applicable to this service or supply. **If there is a discrepancy between a Medical Policy and a member's benefit plan, summary plan description or contract, the benefit plan, summary plan description or contract will govern.**

Coverage

Nonpharmacologic treatment of rosacea, including but not limited to: laser and light therapy, dermabrasion, chemical peels, surgical debulking, and electrosurgery **is considered experimental, investigational and/or unproven.**

NOTE: This policy does not address surgical excision and skin grafting/flap surgery for the treatment of advanced nodular rhinophyma causing a functional impairment (e.g., airway obstruction).

Policy Guidelines

None.

Description

Rosacea is a chronic, inflammatory skin condition without a known cure; the goal of treatment is symptom management. Nonpharmacologic treatments, including laser and light therapy as well as dermabrasion, which are the focus of this medical policy, are proposed for patients who do not want to use or are unresponsive to pharmacologic therapy.

Rosacea

Rosacea is characterized by episodic erythema, edema, papules, pustules, and telangiectasia that occur primarily on the face but also present on the scalp, ears, neck, chest, and back. On occasion, rosacea may affect the eyes. Patients with rosacea tend to flush or blush easily. Because rosacea causes facial swelling and redness, it is easily confused with other skin conditions such as acne, skin allergy, and sunburn.

Rosacea mostly affects adults with fair skin between the ages of 20 and 60 years and is more common in women but often is most severe in men. Rosacea is not life-threatening, but if not treated, it may lead to persistent erythema, telangiectasias, and rhinophyma (hyperplasia and nodular swelling and congestion of the skin of the nose). The etiology and pathogenesis of rosacea are unknown but may result from both genetic and environmental factors. Some theories on the causes of rosacea include blood vessel disorders, chronic *Helicobacter pylori* infection, *Demodex folliculorum* (mites), and immune system disorders.

While the clinical manifestations of rosacea do not usually impact the physical health status of the patient, psychological consequences from the most visually apparent symptoms (i.e., erythema, papules, pustules, telangiectasias) may impact quality of life. Rhinophyma, an end-stage form of chronic acne, has been associated with obstruction of nasal passages and basal cell carcinoma in rare, severe cases. The probability of developing nasal obstruction or basal or squamous cell carcinoma with rosacea is not sufficient to warrant the preventive removal of rhinophymatous tissue.

Treatment

Rosacea treatment can be effective in relieving signs and symptoms. Treatment may include oral and topical antibiotics, isotretinoin, β -blockers, α_2 -adrenergic agonists (e.g., oxymetazoline, clonidine), and anti-inflammatories. Patients are also instructed on various self-care measures such as avoiding skin irritants and dietary items thought to exacerbate acute flare-ups.

Nonpharmacologic therapy has also been tried in patients who cannot tolerate or do not want to use pharmacologic treatments. To reduce visible blood vessels, treat rhinophyma, reduce redness, and improve appearance, various techniques have been used such as laser and light therapy, dermabrasion, chemical peels, surgical debulking, and electrosurgery. Various lasers used include low-powered electrical devices and vascular light lasers to remove telangiectasias, carbon dioxide lasers to remove unwanted tissue from rhinophyma and reshape the nose, and intense pulsed lights that generate multiple wavelengths to treat a broader spectrum of tissue.

Regulatory Status

Several laser and light therapy systems have been cleared for marketing by the U.S. Food and Drug Administration (FDA) through the 510(k) process for various dermatologic indications, including rosacea. For example, rosacea is among the indications for:

- Vbeam laser system (Candela)
- Stellar M22™ laser system (Lumenis)
- excel VT®, excel V®, and xeo® laser systems (Cutera)
- Harmony® XL multi-application platform laser device (Alma Lasers, Israel)
- UV-300 Pulsed Light Therapy System (New Star Lasers)
- CoolTouch® PRIMA Pulsed Light Therapy System (New Star Lasers).

FDA product code: GEX.

Rationale

Medical policies assess the clinical evidence to determine whether the use of technology improves the net health outcome. Broadly defined, health outcomes are the length of life, quality of life, and ability to function including benefits and harms. Every clinical condition has specific outcomes that are important to patients and managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of technology, 2 domains are examined: the relevance, and quality and credibility. To be relevant, studies must represent 1 or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. RCTs are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

Nonpharmacologic Treatment of Rosacea

Clinical Context and Therapy Purpose

The purpose of nonpharmacologic treatments is to provide a treatment option in individuals who have rosacea and do not want to use or are unresponsive to pharmacologic therapies.

The following PICO was used to select literature to inform this policy.

Populations

The relevant population of interest is individuals with rosacea. Rosacea is characterized by episodic erythema, edema, papules and pustules, and telangiectasia that occur primarily on the face. Clinical presentation varies among individuals.

Interventions

The therapies being considered are nonpharmacologic treatments. Nonpharmacologic treatment options include laser and light therapy, dermabrasion, chemical peels, surgical debulking, and electrosurgery. Laser and light therapies are typically used for persistent erythema or telangiectasia. During laser and light therapy, light energy is absorbed by hemoglobin in cutaneous vessels, which leads to vessel heating and coagulation. Lasers vary from low-powered electrical devices and vascular light lasers (for telangiectasias removal) to carbon dioxide lasers and intense pulsed lights that generate multiple wavelengths to treat a broader spectrum of tissue.

Frequency and duration of laser and light therapy sessions vary, from once to twice per month, for several months. Because light-based techniques do not cure rosacea, periodic treatments may be necessary to maintain symptom relief.

Comparators

The comparators of interest are pharmacologic therapies, which include oral and topical antibiotics, isotretinoin, β -blockers, α_2 -adrenergic agonists (e.g., oxymetazoline, clonidine), and anti-inflammatories. The selection of a pharmacological agent is dependent on the clinical features present for an individual (e.g., redness, edema, papules and pustules).

Outcomes

The general outcome of interest is symptom reduction, which may include a change in redness of skin color or change in erythema score or telangiectasia score. Other outcomes of interest include a reduction in pain, subject satisfaction, and improvement in the quality of life.

Outcome measures can be assessed on treatment completion. Because laser and light therapy are not curative, outcomes can be measured months after treatment to assess symptom recurrence.

Study Selection Criteria

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess long-term outcomes and adverse effects, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Consistent with a 'best available evidence approach,' within each category of study design, studies with larger sample sizes and longer durations were sought.
- Studies with duplicative or overlapping populations were excluded.

Systematic Reviews

Table 1 compares the studies included in the systematic reviews. Tables 2 and 3 summarize the characteristics and results of the reviews, respectively.

A meta-analysis by Zhai et al. (2024) compared the efficacy of intense pulsed light and pulsed dye laser for the management of rosacea. (1) Only RCTs comparing these 2 modalities were included, and the primary outcome measures included clearance rates, erythema scores, and pain scores.

A meta-analysis by Chang and Chang (2022) compared the efficacy of pulsed dye laser to intense pulsed light. (2) Only RCTs comparing these 2 modalities were included, and erythema was the only outcome analyzed in meta-analysis.

A meta-analysis by Husein-ElAhmed and Steinhoff (2021) compared the efficacy and tolerability of pulsed dye laser to other laser and light therapies. (3) Both randomized and non-randomized studies were considered for inclusion; background erythema, telangiectasias, pain, and treatment success were analyzed. The studies did not compare interventions with pharmacologic treatments or placebo controls, only pulsed dye laser to other laser and light therapies.

A Cochrane systematic review by van Zuuren et al. (2015) assessed various interventions for rosacea; (4) the same authors updated their systematic review in 2019 with a focus on rosacea phenotypes. (5) In 2019, the authors identified only 7 trials on light and/or laser therapy, and the trials did not compare these interventions with pharmacologic treatments or placebo controls, although 2 studies evaluated laser therapy in combination with pharmacologic therapy. Trial findings on light and/or laser therapy were considered low-quality and were not pooled. The remainder of the RCTs in the review evaluated pharmacologic treatments.

Wat et al. (2014) identified 9 studies on the efficacy of intense pulsed light (IPL) for treating rosacea. (6) Two studies were controlled (left-right comparisons), and the remainder were uncontrolled, including a case report.

The systematic reviews by van Zuuren et al. (2019) and Wat et al. (2014) did not pool study findings on the nonpharmacologic treatment of rosacea. Findings of the published systematic reviews highlight the shortage of RCTs on light and laser therapy for treating rosacea.

Table 1. Comparison of Trials/Studies Included in Systematic Reviews of Nonpharmacologic Treatment of Rosacea

Study	Wat et al. (2014) (6)	van Zuuren et al. (2019) (5)	Husein- ElAhmed and Steinhoff (2021) (3)	Chang and Chang (2022) (2)	Zhai et al. (2024) (1)

West et al. (1998) (7)			●		
Mark et al. (2003) (8)	●				
Taub et al. (2003) (9)	●				
Schroeter et al. (2005) (10)	●				
Karsai et al. (2008) (11)		●			
Papageorgiou et al. (2008) (12)	●				
Neuhaus et al. (2009) (13)	●	●	●		
Lane et al. (2010) (14)	●				
Nymann et al. (2010) (15)		●	●		●
Fabi et al. (2011) (16)	●				
Kassier et al. (2011) (17)	●				
Kim et al. (2011) ^a (18)		●			
Huang et al. (2012) ^a (19)		●			
Tanghetti et al. (2012) (20)			●		●
Alam et al. (2013) (21)		●	●		
Salem et al. (2013) (22)			●		
Friedmann et al. (2014) (23)	●				
Seo et al. (2016) (24)		●	●		
Handler et al. (2017) (25)			●	●	
Kim et al. (2017) (26)			●		
Kwon et al. (2018) (27)			●		

Campos et al. (2019) (28)			●		
Kim et al. (2019) (29)			●	●	●
Gao et al. (2019) (30)					●
Tirico et al. (2020) (31)				●	

^a Study evaluated lasers in combination with other therapies. They are listed for completeness but are not included in the results table below.

Table 2. Systematic Review and Meta-Analysis Characteristics

Study	Dates	Trials	Participants	N (Range)	Design	Duration
Wat et al. (2014) (6)	2003 to 2013	9	Patients with rosacea who received IPL	304 (1 to 102)	2 prospective right-left comparison, 3 OL trials, 3 retrospective, 1 case report	1 to 24 weeks
van Zuuren et al. (2019) (5)	2008 to 2016	7	Patients with rosacea who received laser and light therapies	233 (16 to 60)	RCT	4 to 24 weeks
Husein-ElAhmed and Steinhoff (2021) (3)	1998 to 2019	12	Patients with rosacea who received laser and light therapies	262 (9 to 39)	11 RCTs, 1 prospective right-left comparison	1 to 6 months
Chang and Chang (2022) (2)	2017 to 2020	3	Patients with rosacea who received pulsed dye laser and IPL	29 (5 to 15)	RCT	4 to 12 weeks
Zhai et al. (2024) (1)	Up to 2024	4	Patients with rosacea who received pulsed dye laser and IPL	141 (9 to 39)	RCT	NR

IPL: intense pulsed laser; N: number; NR: not reported; OL: open-label; RCT: randomized controlled trial.

Table 3. Systematic Review & Meta-Analysis Results

Study (Year)	Reduced erythema	Reduced telangiectasia	Reduced blood flow	Visual clearance	VAS pain scores	Adverse events
Wat et al. (2014) (6)						
Total N	300	201	4	60		304
Pooled effect	Seen in 21% to 83% of patients	Seen in 29% to 55% of patients	30% decrease observed in 1 study	Seen in 75% to 87% of patients in 1 study		Included mild itch, edema, bruising, erythema purpura, pain, hyper-pigmentation, and blister
p	p<.05 in 1 study, p<.001 in 1 study	p<.05 in 1 study, p<.001 in 1 study	p<.05 in 1 study	NR		NR
van Zuuren et al. (2019) (5)						
Total N	65	56	NR	40		155
Pooled effect	Low to moderate certainty evidence for IPL, pulsed dye lasers, and Nd:YAG lasers; in 1 study, reduction in erythema index was similar with pulsed dye lasers vs dual wavelength lasers; in 1 study, erythema was reduced with pulsed dye lasers vs Nd:YAG lasers	Low to moderate certainty evidence for IPL, pulsed dye lasers, and Nd:YAG lasers; in 1 study, dual wavelength lasers led to greater improvement vs single wavelength lasers (RR, 4.5); 1 study reported no difference between IPL and pulsed dye laser	NR	Similar number of patients had 75% to 100% response and 50% to 74% response with IPL and long pulsed dye laser		Included purpura, erythema, crusts, hyperpigmentation, vesicles, dryness, itch, tightening, swelling, pain
p	p=.02 in 1 study of pulsed dye	NR	NR	NR		NR

	lasers vs Nd:YAG lasers					
Husein-ElAhmed and Steinhoff (2021) (3)						
Total N	69	NR	NR	148		185
Pooled effect	Pulsed dye lasers vs other laser and light therapies: mean difference, 0.90 (95% CI, -0.99 to 2.79)	Pulsed dye lasers vs other laser and light therapies: RR, 0.54 (95% CI, -0.87 to 1.94)	NR	Treatment success per physician assessment, pulsed dye lasers vs other laser and light therapies: OR, 1.23 (95% CI, 0.74 to 2.04)		Pain, pulsed dye lasers vs other laser and light therapies: mean difference, -0.23 (95% CI, -0.96 to 0.49)
p	p=.35	NR	NR	p=.43		p=.53
Chang and Chang (2022) (2)						
Total N	29	NR	NR	NR		NR
Pooled effect	SMD: -0.112 (95% CI, -0.669 to 0.446)					
p	p=.695					
Zhai et al. (2024) (1)						
Total N	86			141	64	
Pooled Effect				Clearance Rate >50%: incidence of clearance rate >50% was up to 100% in the PDL group and 88.89% in the IPL group (RR, -0.07; 95% CI, -0.19 to 0.05) Clearance Rate >75%:incidence of clearance rate >75% was 66.67%in the PDL group and 77.78% in the IPL group (RR,	SMD= 1.54; (95% CI), 0.08 to 3.00)	

				-0.13; 95% CI, -0.23 to -0.04)		
p	P=.48			Clearance Rate >50%; p=.23 Clearance Rate >75%; p=.007	P=.04	

CI: confidence interval; IPL: intense pulsed light; N: number; Nd:YAG: neodymium-doped yttrium aluminum garnet; NR: not reported; OR: odds ratio; RR: relative risk; SMD: standard mean difference; VAS: visual analog scale.

Randomized Controlled Trials

Several randomized trials evaluating nonpharmacologic treatment for rosacea, almost all of which used split-faced designs, were identified. (11, 13, 21, 28, 29, 31-34) Most compared 2 types of lasers, and only 1 used a placebo control or a pharmacologic treatment as a comparator. Additional RCTs were identified that evaluated the combination of nonpharmacologic and pharmacologic treatments against nonpharmacologic or pharmacologic treatment alone. (35-38) No RCTs evaluating dermabrasion, chemical peels, surgical debulking, or electrosurgery for treating rosacea were identified.

Most studies reported a significant difference in erythema compared to baseline with laser treatments, but no studies found significant differences between laser modalities. For telangiectasia, significant improvements were observed with laser treatments, but only the study by Karsai et al. (2008) reported a significant difference between laser modalities in favor of dual wavelength compared to single wavelength. (11) In the RCT by Campos et al. (2019), the primary outcome of change in Dermatology Life Quality Index was significant compared to baseline after the first ($p<.001$), second ($p=.018$), and third ($p=.001$) treatments. (28) Three studies reported positive findings in subjective measures of patient satisfaction, including patient assessment of change in erythema. (13, 21, 32) Adverse effects in these studies were mild and transient overall. One study reported a significant difference in pain, which was in favor of pulsed dye laser compared to neodymium-doped yttrium aluminum garnet (Nd:YAG) lasers. (21) One RCT reported similar improvements in erythema with pulsed dye laser with topical oxymetazoline compared to topical oxymetazoline alone. (35) A more recent RCT reported greater improvement in erythema with broadband light (intense pulsed light) plus intradermal botulinum toxin compared to broadband light alone. (37)

A summary of key RCT characteristics and results is presented in Tables 4 and 5, respectively. Tables 6 and 7 provide an overview of the relevance and study design/conduct limitations of these RCTs.

Table 4. Summary of Key Randomized Controlled Trial Characteristics

Study	Countries	Sites	Dates	Participants	Description of Interventions	
					Active	Comparator

Karsai et al. (2008) ^a (11)	Germany	1	2006	Patients with nasal telangiectasia with similar vessel densities on both sides and vessel size <0.6 mm	Pulsed dye laser or Nd:YAG laser on 1 side of the face (n=20) Single treatment	Dual wavelength laser on opposite side of the face (n=20)
Neuhaus et al. (2009) ^a (13)	United States	1	NR	Patients age 18 years or older with moderate erythemato-telangiectatic rosacea with background erythema and small vessels (<1 mm) involving the central face	Pulsed dye laser on 1 side of the face (n=22) Pulsed dye laser (n=4) or IPL (n=4) on 1 side of the face 3 treatments separated by 4 weeks each	IPL on opposite side of the face (n=22) No treatment on opposite side of the face (n=8)
Maxwell et al. (2010) ^a (32)	Canada	1	NR	Patients with erythemato-telangiectatic acne rosacea, a personal history of flushing, a family history of rosacea, and rosacea exacerbation by sun, alcohol, and/or spicy food	532 nm long-pulse laser on 1 side of the face (n=11) 6 treatments over 3 months, combined with topical retinaldehyde	Topical retinaldehyde treatment alone on opposite side of the face (n=11)
Alam et al. (2013) ^a (21)	United States	1	NR	Patients age 18 years or older with erythemato-telangiectatic rosacea	Pulsed dye laser on 1 side of the face (n=14)	Nd:YAG laser on opposite side of the face (n=14)

					4 treatments every 3 to 4 weeks	
Campos et al. (2019) ^a (28)	Spain	1	2015	Patients age 18 years or older with erythematotelangiectatic rosacea and no laser treatment within the past year	Pulsed dye laser on 1 side of the face (n=27) 4 treatments every 3 to 4 weeks	Multiplex pulsed dye laser/Nd:YAG laser on opposite side of the face (n=27)
Kim et al. (2019) ^a (29)	Korea	1	NR	Patients with rosacea	Short pulse IPL on 1 side of the face (n=9) 4 treatments every 3 weeks	Pulsed dye laser on opposite side of the face (n=9)
Tirico et al. (2020) ^a (31)	United States	1	2016	Patients age 18 years or older with facial redness and none or mild tan	Short pulse IPL on 1 side of the face (n=5) 2 treatments separated by 4 to 6 weeks	Pulsed dye laser on opposite side of the face (n=5)
Sodha et al. (2021) (35)	United States	1	NR	Patients age 18 years or older with erythematotelangiectatic rosacea	Pulsed dye laser (3 treatments every 4 weeks) plus daily topical oxymetazoline 1% (n=17)	Daily topical oxymetazoline 1% (n=13)

Osman et al. (2022) (36)	Egypt	1	NR	Patients with erythematotelangiectatic or papulopustular rosacea	Pulsed dye laser (4 treatments every 4 weeks) plus daily topical ivermectin 1% (n=15)	Pulsed dye laser alone (n=15)
Tong et al. (2022) ^a (37)	China	1	2021	Patients 14 years or older with rosacea with erythema and flushing as primary symptoms and inadequate response to traditional pharmacologic treatment; no local or systemic pharmacologic treatment within the past 2 weeks	IPL (3 treatments every 4 weeks) plus one-time intradermal botulinum toxin on 1 side of the face (n=22)	IPL plus one-time intradermal saline injection on opposite side of the face (n=22)
Barbarino et al. (2022) ^a (38)	United States	1	NR	Patients 18 to 80 years with moderate-to-severe rosacea including erythema and telangiectasia; no local or systemic therapy within the past 2 weeks	IPL (one treatment) plus the following to right side of face only: phyto-corrective mask application (once per week), phyto-corrective gel (twice daily), topical resveratrol (once daily) (n=10)	IPL (one treatment) alone on opposite side of face (n=10)

Park et al. (2022) ^a (33)	Korea	1	2021	Patients with erythemato-telangiectatic or papulopustular rosacea not treated with antibiotics within the past 4 weeks or with laser treatment within the past 3 months	Long-pulsed alexandrite laser on 1 side of the face (n=23) (4 treatments every 4 weeks)	Pulsed dye laser on opposite side of the face (n=23)
Yang et al. (2023) (34)	China	1	2020-2021	Patients 18 to 65 years with moderate-to-severe rosacea; no phototoxic or photosensitizing drugs within 2 months	ALA-PDT for 3 to 5 sessions (n=20)	Minocycline 100 mg daily for 8 weeks (n=21)

ALA-PDT: 5-aminolevulinic acid photodynamic therapy, IPL: intense pulsed light; Nd:YAG: neodymium-doped yttrium aluminum garnet, NR: not reported.

^a Split face design, yielding an equal number of patients in each treatment group.

Table 5. Summary of Key Randomized Controlled Trial Results/Outcomes

Study (Year)	Change in erythema	Change in telangiectasia	Adverse events
Karsai et al. (2008) (11)	Dual wavelength vs single wavelength		
Percentage, p	NR	>50% vessel clearance: 90% vs 20%, p<.0001	Transient purpura, posttreatment erythema
Neuhaus et al. (2009) (13)	IPL vs pulsed dye laser		
Percentage, p	Malar and alar regions (both treatments): NS Cheek region: IPL vs control, p=.04; Pulsed dye laser vs control, p=.05 All locations: IPL vs pulsed dye laser, NS	Malar and alar region: Pulsed dye laser vs control, both p=.02 IPL vs control, p=.016 and p=.09, respectively IPL vs pulsed dye laser, NS	NR
Maxwell et al. (2010) (32)	Laser vs no laser treatment		
Percentage, p	Mild/moderate improvement: 100%	Mild/moderate improvement: 100%	NR

Alam et al. (2013) (21)	Pulsed dye laser vs Nd:YAG		
Difference (95% CI), p	Pulsed dye laser vs baseline: 8.9% (95% CI, -12.9% to -4.95%), p=.0003 Nd:YAG vs baseline: 2.5% (95% CI, -6.37% to 1.29%), p=.1762 Pulsed dye laser vs Nd:YAG: p=.199	NR	Pain: Worse with Nd:YAG vs pulsed dye laser (p=.0028)
Campos et al. (2019) (28)	Pulsed dye laser vs multiplexed laser		
Difference, p	Erythema index mean change: No difference between treatments (at 3 facial areas), p=.231, p=.674, p=.966, respectively	NR	Adverse effects: 48.1% to 55.6% (pulsed dye laser), purpura most common 14.8% to 33.3% (multiplexed laser), edema most common
Kim et al. (2019) (29)	Short pulse IPL vs pulsed dye laser		
Difference, p	Erythema index mean change: -4.93±1.59 (short pulse IPL) -4.27±1.23 (pulsed dye laser) Difference between treatments: NS	NR	None observed
Tirico et al. (2020) (31)	Short pulse IPL vs pulsed dye laser		
Difference, p	Improvement: 60% vs 45%, NS	NR	Mild pain (mean scores 3.5 to 3.6 for short pulse IPL, mean scores 2.6 to 2.8 for pulsed dye laser)
Sodha et al. (2021) (35)	Pulsed dye laser + oxymetazoline vs oxymetazoline alone		

Difference, p	Clinical Erythema Assessment, change from baseline: Combination: -0.6, -0.7, and -1.2 at 1-, 2-, and 3-months ($p \leq .01$ compared to baseline for all) Oxymetazoline alone: -0.6, -1.2, and -0.9 at 1-, 2-, and 3-months ($p \leq .01$ compared to baseline for all)	NR	Adverse effects: Pulsed dye laser: transient erythema (87%), edema (51%), and purpura (30%) Oxymetazoline (both groups): mild dryness (7%)
Osman et al. (2022) (36)	Pulsed dye laser + ivermectin vs pulsed dye laser alone		
Difference, p	Erythema severity grade significantly reduced compared to baseline in both groups ($p=.005$ for combination, $p=.001$ for pulsed dye laser alone) Difference between treatments: $p = .341$	NR	Mild post-procedural purpura in both groups
Tong et al. (2022) (37)	IPL + intradermal botulinum toxin vs IPL alone		
Difference, p	Erythema index mean change at 3 months: -93.03±42.33 (combination) -66.33±37.53 (IPL alone) Difference between treatments: $p < .05$	NR	Mild erythema and pain at injection site
Barbarino et al. (2022) (38)	Phyto-corrective therapy + IPL vs IPL alone		
Difference, p	Not individually reported Efficacy represented by physician-assessed global aesthetic improvement scale at 3 months relative to baseline: Combination: 50%, 20%, and 30% improved, much	Not individually reported	NR

	improved, and very much improved, respectively IPL alone: 10%, 60%, and 20% improved, much improved, and very much improved, respectively		
Park et al. (2022) (33)	Long-pulsed alexandrite laser plus pulsed dye laser vs pulsed dye laser alone		
Difference, p	Erythema index mean change at 3 months after last treatment: -20.1%±15.4% (combination) -23.0%±18.7% (pulsed dye laser alone) Difference between treatments: p=.325	NR	Transient erythema or swelling
Yang et al. (2023) (34)	Reduction in lesions (median, IQR)	Treatment success (n, %)	RosaQol change (median, IQR)
Difference, p	ALA-PDT: 19 (12 to 36) Minocycline: 22 (12 to 40) p=.75	ALA-PDT: 16 (80%) Minocycline: 17 (81%) p=1	ALA-PDT: 0.48 (0.19 to 1.22) Minocycline: 0.53 (-0.27 to 1.57) p=.64

ALA-PDT: 5-aminolevulinic acid photodynamic therapy, CI: confidence interval; IPL: intense pulsed laser; IQR: interquartile range; Nd:YAG: neodymium-doped yttrium aluminum garnet; NR: not reported; NS: not significant; RosaQol: Rosacea-specific Quality-of-Life.

Table 6. Study Relevance Limitations

Study	Population^a	Intervention^b	Comparator^c	Outcomes^d	Follow-Up^e
Karsai et al. (2008) (11)			2 - no comparison to established pharmacologic treatment group alone	5 - clinically significant difference not prespecified	1 - only 1 treatment
Neuhaus et al. (2009) (13)			2 - no comparison to established pharmacologic treatment group alone	3 - no mention of harms 5 - clinically significant difference not prespecified	

Maxwell et al. (2010) (32)				3 - no mention of harms 4 - major outcomes were patient-related subjective improvements 5 - clinically significant difference not prespecified	
Alam et al. (2013) (21)			2 - no comparison to established pharmacologic treatment group alone	5 - clinically significant difference not prespecified	
Campos et al. (2019) (28)			2 - no comparison to established pharmacologic treatment group alone	5 - clinically significant difference not prespecified	
Kim et al. (2019) (29)			2 - no comparison to established pharmacologic treatment group alone	5 - clinically significant difference not prespecified	
Tirico et al. (2020) (31)				5 - clinically significant difference not prespecified	
Sodha et al. (2021) (35)			2 - missing inclusion of a laser-based treatment group only	5 - clinically significant difference not prespecified	
Osman et al. (2022) (36)			2 - no comparison to established pharmacologic	5 - clinically significant difference not prespecified	

			treatment group alone		
Tong et al. (2022) (37)			2 - no comparison to established pharmacologic treatment group alone	5 - clinically significant difference not prespecified	
Barbarino et al. (2022) (38)	4 - only enrolled women	1 - details of intervention formulations and dosing unclear	2 - only a single laser therapy treatment administered; no comparison to established pharmacologic treatment group alone	1 - erythema, telangiectasia, and other disease outcomes not individually reported 3 - no reporting of harms 4 - invalid patient-reported outcomes 5 - clinically significant difference not specified 6 - clinically significant difference not supported	
Park et al. (2022) (33)			2 - no comparison to established pharmacologic treatment group alone	3 - incomplete reporting of harms 5 - clinically significant difference not prespecified	
Yang et al. (2023) (34)					

ALA-PDT: 5-aminolevulinic acid photodynamic therapy

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Population key: 1. Intended use population unclear; 2. Clinical context for treatment is unclear; 3. Study population unclear; 4. Study population not representative of intended use; 5. Study population is subpopulation of intended use.

^b Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator.

^c Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively.

^d Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. Not CONSORT reporting of harms; 4. Not established and validated measurements; 5. Clinically significant difference not prespecified; 6. Clinically significant difference not supported.

^e Follow-up key: 1. Not sufficient duration for benefits; 2. Not sufficient duration for harms.

Table 7. Study Design and Conduct Limitations

Study	Allocation^a	Blinding^b	Selective Reporting^c	Follow-Up^d	Power^e	Statistical^f
Karsai et al. (2008) (11)		1-no mention of patient blinding			1-no mention of power	3-p-value for primary efficacy comparison not reported
Neuhaus et al. (2009) (13)		1-no mention of patient blinding			1-no mention of power	
Maxwell et al. (2010) (32)		1-no mention of patient blinding 2-most outcomes were patient-related improvements and patients were not blinded		6 - only reported results for patients that completed the study	1-no mention of power	4-treatments were not statistically compared
Alam et al. (2013) (21)				6 - only reported results for patients that completed the study		
Campos et al. (2019) (28)				6 - only reported results for patients that completed the study	1-no mention of power	

Kim et al. (2019) (29)		1-no mention of patient blinding 2-no mention of blinding				
Tirico et al. (2020) (31)				6 - only reported results for patients that completed the study	1-no mention of power	3-p-value for primary efficacy comparisons not reported
Sodha et al. (2021) (35)		1-no mention of patient blinding 2-some outcomes were patient-rated improvements , and patients were not blinded			2-power not reported for primary outcome; authors noted adequate power not achieved due to closure of the clinic due to COVID- 19	4-change in erythema not compared between treatment arms
Osman et al. (2022) (36)		1 - no mention of patient blinding	1 - not registered		1 - no mention of power	1 - test used to compare between arms unclear 2 - unclear if appropriate test used for multiple observations
Tong et al. (2022) (37)		1 - no mention of patient blinding			1 - no mention of power	
Barbarino et al.		1 - no mention of blinding	1 - not registered		1 - no inferential	3 - no inferential

(2022) (38)		3 - outcome assessed by treating physician	2 - evaluation of reduction in procedure-related adverse events with intervention stated in study aims, but no safety results reported 3 - senior author is journal's editor-in-chief, third author is on journal's advisory committee		statistical analysis	statistical analysis 4 - no inferential statistical analysis
Park et al. (2022) (33)				2 - details of handling data for dropout cases not reported 6 - dropout cases appear to be excluded from analysis, procedure not detailed	1 - no mention of power	

Yang et al. (2023) (34)		1 - patients not blind				
-------------------------------	--	---------------------------	--	--	--	--

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

^a Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias.

^b Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician.

^c Selective reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

^d Follow-up key: 1. High loss to follow up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials).

^e Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference.

^f Statistical key: 1. Test is not appropriate for outcome type: a) continuous; b) binary; c) time to event; 2. Test is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p-values not reported; 4. Comparative treatment effects not calculated.

Summary of Evidence

For individuals who have rosacea who receive nonpharmacologic treatment (e.g., laser therapy, light therapy, dermabrasion), the evidence includes systematic reviews and several small, randomized, split-face design trials. Relevant outcomes are symptoms, change in disease status, and treatment-related morbidity. The systematic reviews reported favorable effects on erythema and telangiectasia with several laser types, including intense pulsed light (IPL), pulsed dye lasers, and neodymium-doped yttrium aluminum garnet (Nd:YAG) lasers. However, the systematic reviews did not pool results from individual studies and the studies differed in the specific lasers being compared. Overall, the systematic review results were insufficient to establish whether any laser type is more effective and safer than others. The randomized controlled trials (RCTs) evaluated laser and light therapy. One RCT compared combination laser and pharmacologic therapy with pharmacologic therapy alone and 2 RCTs compared combination laser and pharmacologic therapy with laser therapy alone, but the lack of an arm evaluating laser therapy alone against established pharmacologic therapy does not allow a direct assessment on the efficacy of laser or light treatment compared with alternative treatments. No trials assessing other nonpharmacologic treatments were identified. There is a need for RCTs that compare nonpharmacologic treatments with placebo controls and with pharmacologic treatments. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Practice Guidelines and Position Statements

American Acne and Rosacea Society

In 2014, the American Acne and Rosacea Society issued consensus recommendations on the management of rosacea. (39) The Society stated that lasers and intense pulsed light (IPL) devices could improve certain clinical manifestations of rosacea that have not responded to

medical therapy. The recommendations indicated that these therapies would have to be repeated intermittently to sustain improvement.

In 2020, the American Acne and Rosacea Society issued updated consensus recommendations on the management of rosacea. (40) The Society's consensus recommendation on rosacea management correlated with clinical manifestations observed at the time of presentation is summarized in Table 8.

Table 8. Recommendations on Use of Lasers and Intense Pulsed Light Devices for the Management of Rosacea

Condition	Recommendation	Grade ^a
Persistent central facial erythema without papulopustular lesions	Intense pulsed light (IPL), potassium titanyl phosphate crystal laser (KTP), or pulsed-dye laser	B
Diffuse central facial erythema with papulopustular (PP) lesions	“While the data on the use of IPL, KTP or pulsed-dye laser are limited for PP lesions, these options are useful to treat erythema”	NR
Granulomatous rosacea	• Intense pulsed dye laser • “No current standard of treatment; limited data based on case reports”	C
Phymatous Rosacea	• “Surgical therapy for fully developed phymatous changed (carbon dioxide laser, erbium-doped [YAG] laser, electrosurgery, dermabrasion)” • “Treatment selection dependent on stage of development (early or fibrotic) and extent of inflammation (active or burnt out)”	C

IPL: intense pulsed light, YAG: yttrium aluminum garnet; NR: not reported.

^a Grade A: Criteria not described in recommendation; Grade B: Systematic review/meta-analysis of lower-quality clinical trials or studies with limitations and inconsistent findings; lower-quality clinical trial; Grade C: Consensus guidelines; usual practice, expert opinion, case series—limited trial data

National Rosacea Society

In 2019, the National Rosacea Society Executive Committee published an expert consensus document on management options for rosacea. (41) This document endorses treatment goals of an Investigator Global Assessment score of 0 and normalization of skin tone and color due to the notable impact of rosacea on patient quality of life. Light devices are discussed as treatment options along with medications, skin care, and lifestyle interventions. Based on weak evidence, IPL, pulsed dye lasers, and potassium titanyl phosphate lasers are listed as moderately effective treatment options for persistent erythema, particularly due to telangiectasia. Both IPL and potassium titanyl phosphate are described as having at least some efficacy for flushing. Nonpharmacologic interventions that are listed as more highly effective treatment options for non-inflamed phymas (based on weak evidence) include carbon dioxide lasers, erbium lasers, cold steel, electrosurgery, and radiofrequency; these same interventions are listed for use in

combination with other treatment modalities for inflammatory phymas. Carbon dioxide lasers, erbium lasers, cold steel, electrosurgery, and radiofrequency carry a risk of post-inflammatory hyperpigmentation and should only be provided by appropriately trained individuals.

Rosacea Consensus Panel

In 2017, the Rosacea Consensus panel, comprised of international experts including representatives from the U.S., published recommendations for rosacea treatment. (42) The panel agreed that treatments should be based on phenotype. IPL and pulsed dye laser were recommended for persistent erythema, but not for transient erythema. IPL and lasers were also recommended for telangiectasia rosacea.

The panel updated their recommendations on rosacea treatment in 2019, agreeing that lasers were recommended for persistent centrofacial erythema. (43) They also noted that “use of IPL and vascular lasers in darker skin phototypes requires consideration by a healthcare provider with experience..., as it can result in dyspigmentation.” The panel also acknowledged that combining treatments could benefit patients with more severe rosacea and multiple rosacea features; however, “there remains an ongoing need for more studies to support combination treatment use in rosacea.”

Ongoing and Unpublished Clinical Trials

No ongoing or unpublished trials were identified in a search of clinicaltrials.gov in November 2025.

Coding

Procedure codes on Medical Policy documents are included **only** as a general reference tool for each policy. **They may not be all-inclusive.**

The presence or absence of procedure, service, supply, or device codes in a Medical Policy document has no relevance for determination of benefit coverage for members or reimbursement for providers. **Only the written coverage position in a Medical Policy should be used for such determinations.**

Benefit coverage determinations based on written Medical Policy coverage positions must include review of the member’s benefit contract or Summary Plan Description (SPD) for defined coverage vs. non-coverage, benefit exclusions, and benefit limitations such as dollar or duration caps.

CPT Codes	15260, 15261, 15780, 15781, 15782, 15783, 15788, 15789, 15792, 15793, 17000, 17003, 17004, 17106, 17107, 17108, 30117, 30118, 30120
HCPCS Codes	None

*Current Procedural Terminology (CPT®) ©2025 American Medical Association: Chicago, IL.

References

1. Zhai Q, Cheng S, Liu R, et al. Meta-Analysis of the Efficacy of Intense Pulsed Light and Pulsed-Dye Laser Therapy in the Management of Rosacea. J Cosmet Dermatol. Dec 2024; 23(12):3821-3827. PMID 39240125

2. Chang HC, Chang YS. Pulsed dye laser versus intense pulsed light for facial erythema of rosacea: a systematic review and meta-analysis. *J Dermatolog Treat.* Jun 2022; 33(4):2394-2396. PMID 34291712
3. Husein-ElAhmed H, Steinhoff M. Light-based therapies in the management of rosacea: a systematic review with meta-analysis. *Int J Dermatol.* Feb 2022; 61(2):216-225. PMID 34089264
4. van Zuuren EJ, Fedorowicz Z, Carter B, et al. Interventions for rosacea. *Cochrane Database Syst Rev.* Apr 28 2015; 2015(4):CD003262. PMID 25919144
5. van Zuuren EJ, Fedorowicz Z, Tan J, et al. Interventions for rosacea based on the phenotype approach: an updated systematic review including GRADE assessments. *Br J Dermatol.* Jul 2019; 181(1):65-79. PMID 30585305
6. Wat H, Wu DC, Rao J, et al. Application of intense pulsed light in the treatment of dermatologic disease: a systematic review. *Dermatol Surg.* Apr 2014; 40(4):359-377. PMID 24495252
7. West TB, Alster TS. Comparison of the long-pulse dye (590-595 nm) and KTP (532 nm) lasers in the treatment of facial and leg telangiectasias. *Dermatol Surg.* Feb 1998; 24(2):221-226. PMID 9491116
8. Mark KA, Sparacio RM, Voigt A, et al. Objective and quantitative improvement of rosacea-associated erythema after intense pulsed light treatment. *Dermatol Surg.* Jun 2003; 29(6):600-604. PMID 12786702
9. Taub AF. Treatment of rosacea with intense pulsed light. *J Drugs Dermatol.* Jun 2003; 2(3):254-259. PMID 12848109
10. Schroeter CA, Haaf-von Below S, Neumann HA. Effective treatment of rosacea using intense pulsed light systems. *Dermatol Surg.* Oct 2005; 31(10):1285-1289. PMID 16188180
11. Karsai S, Roos S, Raulin C. Treatment of facial telangiectasia using a dual-wavelength laser system (595 and 1,064 nm): a randomized controlled trial with blinded response evaluation. *Dermatol Surg.* May 2008; 34(5):702-708. PMID 18318728
12. Papageorgiou P, Clayton W, Norwood S, et al. Treatment of rosacea with intense pulsed light: significant improvement and long-lasting results. *Br J Dermatol.* Sep 2008; 159(3):628-632. PMID 18565174
13. Neuhaus IM, Zane LT, Tope WD. Comparative efficacy of nonpurpuragenic pulsed dye laser and intense pulsed light for erythematotelangiectatic rosacea. *Dermatol Surg.* Jun 2009; 35(6):920-928. PMID 19397667
14. Lane JE, Khachemoune A. Use of intense pulsed light to treat refractory granulomatous rosacea. *Dermatol Surg.* Apr 2010; 36(4):571-573. PMID 20402938
15. Nymann P, Hedelund L, Haedersdal M. Long-pulsed dye laser vs. intense pulsed light for the treatment of facial telangiectasias: a randomized controlled trial. *J Eur Acad Dermatol Venereol.* Feb 2010; 24(2):143-146. PMID 20205349
16. Fabi S, Peterson J, Goldman M. Combination 15% azelaic acid gel and intense pulse light therapy for mild to moderate rosacea. *Lasers Surg Med.* 2011; 43:9689.
17. Kassir R, Kolluru A, Kassir M. Intense pulsed light for the treatment of rosacea and telangiectasias. *J Cosmet Laser Ther.* Oct 2011; 13(5):216-222. PMID 21848421
18. Kim TG, Roh HJ, Cho SB, et al. Enhancing effect of pretreatment with topical niacin in the treatment of rosacea-associated erythema by 585-nm pulsed dye laser in Koreans: a

- randomized, prospective, split-face trial. *Br J Dermatol*. Mar 2011; 164(3):573-579. PMID 21143465
19. Huang YE, Li XL, Li TJ. [Clinical research of topical tacrolimus ointment combined with 585 nm pulsed dye laser in the treatment of rosacea]. *J Clinical Dermatol*. 2012; 41:3089.
 20. Tanghetti EA. Split-face randomized treatment of facial telangiectasia comparing pulsed dye laser and an intense pulsed light handpiece. *Lasers Surg Med*. Feb 2012; 44(2):97-102. PMID 22180317
 21. Alam M, Voravutinon N, Warycha M, et al. Comparative effectiveness of nonpurpuragenic 595-nm pulsed dye laser and microsecond 1064-nm neodymium:yttrium-aluminum-garnet laser for treatment of diffuse facial erythema: A double-blind randomized controlled trial. *J Am Acad Dermatol*. Sep 2013; 69(3):438-443. PMID 23688651
 22. Salem SA, Abdel Fattah NS, Tantawy SM, et al. Neodymium-yttrium aluminum garnet laser versus pulsed dye laser in erythemato-telangiectatic rosacea: comparison of clinical efficacy and effect on cutaneous substance (P) expression. *J Cosmet Dermatol*. Sep 2013; 12(3):187-194. PMID 23992160
 23. Friedmann DP, Goldman MP, Fabi SG, et al. The effect of multiple sequential light sources to activate aminolevulinic Acid in the treatment of actinic keratoses: a retrospective study. *J Clin Aesthet Dermatol*. Sep 2014; 7(9):20-25. PMID 25276272
 24. Seo HM, Kim JI, Kim HS, et al. Prospective Comparison of Dual Wavelength Long-Pulsed 755-nm Alexandrite/1,064-nm Neodymium:Yttrium-Aluminum-Garnet Laser versus 585-nm Pulsed Dye Laser Treatment for Rosacea. *Ann Dermatol*. Oct 2016; 28(5):607-614. PMID 27746641
 25. Handler MZ, Bloom BS, Goldberg DJ. IPL vs PDL in treatment of facial erythema: A split-face study. *J Cosmet Dermatol*. Dec 2017; 16(4):450-453. PMID 28752575
 26. Kim SJ, Lee Y, Seo YJ, et al. Comparative Efficacy of Radiofrequency and Pulsed Dye Laser in the Treatment of Rosacea. *Dermatol Surg*. Feb 2017; 43(2):204-209. PMID 27893539
 27. Kwon WJ, Park BW, Cho EB, et al. Comparison of efficacy between long-pulsed Nd:YAG laser and pulsed dye laser to treat rosacea-associated nasal telangiectasia. *J Cosmet Laser Ther*. Oct 2018; 20(5):260-264. PMID 29388843
 28. Campos MA, Sousa AC, Varela P, et al. Comparative effectiveness of purpuragenic 595 nm pulsed dye laser versus sequential emission of 595 nm pulsed dye laser and 1,064 nm Nd:YAG laser: a double-blind randomized controlled study. *Acta Dermatovenereol Alp Pannonica Adriat*. Mar 2019; 28(1):1-5. PMID 30901061
 29. Kim BY, Moon HR, Ryu HJ. Comparative efficacy of short-pulsed intense pulsed light and pulsed dye laser to treat rosacea. *J Cosmet Laser Ther*. Aug 2019; 21(5):291-296. PMID 30285506
 30. Gao L, Qu H, Gao N, et al. A retrospective analysis for facial telangiectasia treatment using pulsed dye laser and intense pulsed light configured with different wavelength bands. *J Cosmet Dermatol*. Jan 2020; 19(1):88-92. PMID 31638320
 31. Tirico MCCC, Jensen D, Green C, et al. Short pulse intense pulsed light versus pulsed dye laser for the treatment of facial redness. *J Cosmet Laser Ther*. Feb 17 2020; 22(2):60-64. PMID 32041440

32. Maxwell EL, Ellis DA, Manis H. Acne rosacea: effectiveness of 532 nm laser on the cosmetic appearance of the skin. *J Otolaryngol Head Neck Surg*. Jun 2010; 39(3):292-296. PMID 20470675
33. Park S, Lee JH, Kang E, et al. A randomized split-face comparative study of long-pulsed alexandrite plus low-fluence Nd:YAG laser versus pulsed-dye laser in the treatment of rosacea. *Lasers Surg Med*. Nov 2022; 54(9):1217-1225. PMID 36183378
34. Yang J, Liu X, Cao Y, et al. 5-Aminolevulinic acid photodynamic therapy versus minocycline for moderate-to-severe rosacea: A single-center, randomized, evaluator-blind controlled study. *J Am Acad Dermatol*. Oct 2023; 89(4):711-718. PMID 37356626
35. Sodha P, Suggs A, Munavalli GS, et al. A Randomized Controlled Pilot Study: Combined 595-nm Pulsed Dye Laser Treatment and Oxymetazoline Hydrochloride Topical Cream Superior to Oxymetazoline Hydrochloride Cream for Erythematotelangiectatic Rosacea. *Lasers Surg Med*. Dec 2021; 53(10):1307-1315. PMID 34233378
36. Osman M, Shokeir HA, Hassan AM, et al. Pulsed dye laser alone versus its combination with topical ivermectin 1% in treatment of Rosacea: a randomized comparative study. *J Dermatolog Treat*. Feb 2022; 33(1):184-190. PMID 32141785
37. Tong Y, Luo W, Gao Y, et al. A randomized, controlled, split-face study of botulinum toxin and broadband light for the treatment of erythematotelangiectatic rosacea. *Dermatol Ther*. May 2022; 35(5):e15395. PMID 35187781
38. Barbarino SC, Bucay VW, Cohen JL, et al. Integrative skincare trial of intense pulsed light followed by the phyto-corrective mask, phyto-corrective gel, and resveratrol BE for decreasing post-procedure downtime and improving procedure outcomes in patients with rosacea. *J Cosmet Dermatol*. Sep 2022; 21(9):3759-3767. PMID 35765796
39. Tanghetti E, Del Rosso JQ, Thiboutot D, et al. Consensus recommendations from the American Acne & Rosacea Society on the management of rosacea, part 4: a status report on physical modalities and devices. *Cutis*. Feb 2014; 93(2):71-76. PMID 24605343
40. Del Rosso JQ, Tanghetti E, Webster G, et al. Update on the Management of Rosacea from the American Acne & Rosacea Society (AARS). *J Clin Aesthet Dermatol*. June 2020; 13(6 Suppl):S17-S24. PMID 33282106
41. Thiboutot D, Anderson R, Cook-Bolden F, et al. Standard management options for rosacea: The 2019 update by the National Rosacea Society Expert Committee. *J Am Acad Dermatol*. Jun 2020; 82(6):1501-1510. PMID 32035944
42. Schaller M, Almeida LM, Bewley A, et al. Rosacea treatment update: recommendations from the global ROSacea Consensus (ROSCO) panel. *Br J Dermatol*. Feb 2017; 176(2):465-471. PMID 27861741
43. Schaller M, Almeida LMC, Bewley A, et al. Recommendations for rosacea diagnosis, classification and management: update from the global ROSacea Consensus 2019 panel. *Br J Dermatol*. May 2020; 182(5):1269-1276. PMID 31392722

Centers for Medicare and Medicaid Services (CMS)

The information contained in this section is for informational purposes only. HCSC makes no representation as to the accuracy of this information. It is not to be used for claims adjudication for HCSC Plans.

The Centers for Medicare and Medicaid Services (CMS) does not have a national Medicare coverage position. Coverage may be subject to local carrier discretion.

A national coverage position for Medicare may have been developed since this medical policy document was written. See Medicare's National Coverage at <<https://www.cms.hhs.gov>>.

Policy History/Revision	
Date	Description of Change
05/01/2026	Document updated. Coverage unchanged. References 1 and 30 added; others updated.
09/01/2025	Document updated with literature review. The following changes were made to Coverage: 1) Removed criteria specific to surgical excision and skin grafting/flap surgery for the treatment of advanced nodular rhinophyma; 2) Added "surgical debulking" to list of examples of nonpharmacologic treatments in experimental, investigational and/or unproven statement; and 3) Added "NOTE: This policy does not address surgical excision and skin grafting/flap surgery for the treatment of advanced nodular rhinophyma causing a functional impairment (e.g., airway obstruction)." Added reference 32; another removed.
10/15/2024	Document updated with literature review. Coverage unchanged. References 1, 31-32, 34-36 and 46 added; others updated.
07/01/2023	Reviewed. No changes.
04/15/2022	Document updated with literature review. Coverage unchanged. References 1, 3, 5-8, 10, 12-18, 20, 22-23, 25-27, 29-30, 36, and 39 added.
02/15/2021	Reviewed. No changes.
09/15/2020	Document updated with literature review. Coverage unchanged. References 13 and 15 added.
10/01/2019	Reviewed. No change(s).
03/01/2019	Document updated with literature review. The following changes were made in Coverage: 1) Added medically necessary criteria for the surgical treatment of rhinophyma when there is a functional impairment 2) Moved and expanded the cosmetic statement previously on 716.001 Cosmetic and Reconstructive Services to state "surgical treatment or surgical reshaping of the nose for rhinophyma is considered cosmetic for all other indications or when the above criterion is not met" 3) Removed surgical debulking from the experimental, investigational and/or unproven statement. Added references 9-14.
04/15/2017	Reviewed. No changes.
05/01/2016	Document updated with literature review. Coverage unchanged.

05/15/2015	Reviewed. No changes.
05/15/2014	Document updated with literature review. Coverage unchanged.
04/15/2013	Document updated with literature review. Coverage unchanged.
07/01/2011	Document updated with literature review. References added. Coverage unchanged.
01/01/2009	New medical document.